

Datasheet: MCA1799 BATCH NUMBER 168458

Description:	MOUSE ANTI HUMAN ESTROGEN RECEPTOR ALPHA
Specificity:	ESTROGEN RECEPTOR ALPHA
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	6F11
Isotype:	IgG1
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen (1)	•			1/40 - 1/60
Immunohistology - Paraffin (2)	-			1/40 - 1/80
ELISA				
Immunoprecipitation				
Western Blotting	•			1/50 - 1/100

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

- (1)The use of Zamboni's fixative is recommended for best results
- (2)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Product Form	Tissue culture supernatant - liquid
Preservative Stabilisers	0.09% sodium azide (NaN ₃)
lmmunogen	Recombinant human estrogen receptor (alpha form).

External Database Links	UniProt:
	P03372 Related reagents
	Entrez Gene:
	2099 ESR1 Related reagents
Synonyms	ESR, NR3A1
RRID	AB_2102070
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse p3-NS1-Ag4-1 myeloma cell line.
Specificity	Mouse anti Human estrogen alpha antibody, clone 6F11 recognizes the human estrogen receptor alpha chain (ER α), also known as the estradiol receptor or nuclear receptor subfamily 3 group A member 1. ER α is a ~65 kDa steroid hormone receptor containing an N-terminal (<u>AF-1</u>) ligand independent transactivation domain, a DNA binding domain and a C-terminal ligand binding domain which overlaps with an (<u>AF-2</u>) domain. ER α binds to DNA as a homodimer (<u>Klinge 2001</u>) and can also form heterodimers with Estrogen receptor beta.
	The detection of estrogen (ER) and progesterone (PR) receptors using immunohistochemical staining of formalin fixed, paraffin embedded (FFPE) tissue, has gradually replaced ligand binding assays, to become the most common method for the determination of the ER/PR status of breast tumors (Yaziji et al. 2008). Approximately 75% to 80% of breast tumors have estrogen and/or progesterone receptors, and the presence of these receptors helps determine both the patients prognosis and the effectiveness of hormonal therapy (Bhargava et al. 2012).
	Mouse anti human estrogen alpha antibody, clone 6F11 has been used successfully for identification of ER α on breast cancer cell lysates using Western blotting and for the immunohistochemical detection of ER α in breast cancer tissues (Ambroise et al. 2011).
Histology Positive Control Tissue	Human breast carcinoma
References	 Bevitt, D.J. <i>et al.</i> (1997) New monoclonal antibodies to oestrogen and progesterone receptors effective for paraffin section immunohistochemistry. <u>J Pathol. 183 (2): 228-32.</u> Jongen, V. <i>et al.</i> (2009) Expression of estrogen receptor-alpha and -beta and progesterone receptor-A and -B in a large cohort of patients with endometrioid endometrial cancer. <u>Gynecol Oncol. 112: 537-42.</u> Ambroise, M. <i>et al.</i> (2011) Immunohistochemical profile of breast cancer patients at a tertiary care hospital in South India. <u>Asian Pac J Cancer Prev. 12: 625-9.</u> Krishnaswamy, U. <i>et al.</i> (2013) Correlation of Her-2 neu over-expression with clinico pathological features of carcinoma breast <u>Apollo Medicine. 10 (4): 313-7.</u> Droog, M. <i>et al.</i> (2017) Estrogen receptor α wields treatment-specific enhancers

between morphologically similar endometrial tumors. Proc Natl Acad Sci U S A. 114 (8):

E1316-E1325.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee Guaranteed until date of expiry. Please see product label.

Health And Safety Material Safety Datasheet documentation #10053 available at:

Information https://www.bio-rad-antibodies.com/SDS/MCA1799

10053

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) HRP
Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...)

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M418601:230427'

Printed on 22 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint