Datasheet: MCA1815 BATCH NUMBER 167814

Description:	MOUSE ANTI HUMAN CD68
Specificity:	CD68
Other names:	MACROSIALIN
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	514H12
lsotype:	lgG2a
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin (1)	•			1/40 - 1/80
ELISA			•	
Immunoprecipitation				
Western Blotting		•		
Immunofluorescence	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Tris/EDTA buffer pH 9.0 is recommended for this purpose. *N.B.* The epitope recognised by clone 514H12 is sensitive to peroxide. It is therefore recommended that any peroxidase-blocking steps be performed after incubation with this antibody.

Target Species	Human
Product Form	Tissue culture supernatant - liquid

Preservative Stabilisers	0.09% sodium azide (NaN ₃)			
Immunogen	Fusion protein corresponding to external domain of human CD68.			
External Database Links	UniProt: <u>P34810</u> <u>Related reagents</u> Entrez Gene: <u>968</u> CD68 <u>Related reagents</u>			
RRID	AB_322866			
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse p3-NS1-Ag4-1 myeloma cell line			
Specificity	Mouse anti Human CD68 antibody, clone 514H12 recognizes the human CD68 cell surface antigen, a ~110 kDa glycoprotein primarily expressed by macrophages and monocytes.			
Histology Positive Control Tissue	Human tonsil			
References	 da Costa, C.E. <i>et al.</i> (2005) Presence of osteoclast-like multinucleated giant cells in the bone and nonostotic lesions of Langerhans cell histiocytosis. J Exp Med. 201 (5): 687-93. Rodriguez-Agudo, D. <i>et al.</i> (2006) Localization of StarD5 cholesterol binding protein. J Lipid Res. 47: 1168-75. Paulmyer-Lacroix, O. <i>et al.</i> (2006) Expression of adrenomedullin in adipose tissue of lean and obese women. Eur J Endocrinol. 155: 177-85. Wang, X. <i>et al.</i> (2006) Monocyte/macrophage and T-cell infiltrates in peritoneum of patients with ovarian cancer or benign pelvic disease. J Transl Med. 4: 30. Angel, C.E. <i>et al.</i> (2007) CD14+ antigen-presenting cells in human dermis are less mature than their CD1a+ counterparts. Int Immunol. 19: 1271-9. Achard, V. <i>et al.</i> (2007) Renin receptor expression in human adipose tissue. Am J Physiol Regul Integr Comp Physiol. 292: R274-82. Hever, A. <i>et al.</i> (2007) Expression of adrenomedullin in human epicardial adipose tissue: role of coronary status. Am J Physiol Endocrinol Metab. 293: E1443-50. Kaibara, N. <i>et al.</i> (2008) Comparative histopathological analysis between tenosynovitis and joint synovitis in rheumatoid arthritis. Histopathology. 52: 856-64. Moskovszky, L. <i>et al.</i> (2011) Use of macrophages to target therapeutic adenovirus to human prostate tumors. Cancer Res. 71 (5): 1805-15. de Vos van Steenwijk PJ <i>et al.</i> (2013) Tumor-infiltrating CD14-positive myeloid cells and CD8-positive T-cells prolong survival in patients with cervical carcinoma. Int J Cancer. 			

<u>133 (12): 2884-94.</u>
13. Zhang, W. et al. (2013) Myeloid clusters are associated with a pro-metastatic
environment and poor prognosis in smoking-related early stage non-small cell lung
cancer. PLoS One. 8 (5): e65121.
14. Toh, M.L. <i>et al.</i> (2014) Bone- and cartilage-protective effects of a monoclonal antibody
against colony-stimulating factor 1 receptor in experimental arthritis. Arthritis Rheumatol.
<u>66 (11): 2989-3000.</u>
15. Schirmer, L. <i>et al.</i> (2014) Differential loss of KIR4.1 immunoreactivity in multiple
sclerosis lesions. Ann Neurol. 75 (6): 810-28.
16. Bobosha, K. <i>et al.</i> (2014) T-cell regulation in lepromatous leprosy. <u>PLoS Negl Trop</u>
Dis. 8 (4): e2773.
17. Quispel, W.T. <i>et al.</i> (2016) Tertiary lymphoid structures are confined to patients
presenting with unifocal Langerhans Cell Histiocytosis. <u>Oncoimmunology. 5 (8): e1164364.</u>
18. Schrevel, M. <i>et al.</i> (2017) Autocrine expression of the epidermal growth factor receptor
ligand heparin-binding EGF-like growth factor in cervical cancer. Int J Oncol. 50 (6):
1947-54.
19. van Poppel, P.C.M. <i>et al.</i> (2018) Inflammation in the subcutaneous adipose tissue
does not attenuate endothelial function in subjects with diabetes mellitus and subjects with
dyslipidaemia and hypertension: A cross-sectional study. Endocrinol Diabetes Metab. 1
(<u>3</u>): e00020.
20. Yuan, A. <i>et al.</i> (2020) Histologic analysis of medication-related osteonecrosis of the
jaw compared with antiresorptive-exposed bone and other infectious, inflammatory, and
necrotic jaw diseases. Oral Surg Oral Med Oral Pathol Oral Radiol. 129 (2): 133-40.
21. Durán, D. <i>et al.</i> (2020) Histological and Immunohistochemical Study of Wounds in
Sheep Skin in Maggot Therapy by Using <i>Protophormia terraenovae.</i> (Diptera:
<i>Calliphoridae</i> .) Larvae. <u>J Med Entomol. 57 (2): 369-76.</u>
22. Nowak-Sliwinska, P. et al. (2022) Proinflammatory activity of VEGF-targeted treatment
through reversal of tumor endothelial cell anergy. <u>Angiogenesis</u> . <u>Dec 02 [Epub ahead of</u>
print].
23. Cossins, B.C. <i>et al.</i> (2023) Sex-specific association between adipose tissue
inflammation and vascular and metabolic complications of obesity. <u>J Clin Endocrinol</u>
Metab. dgad193.
24. Perga, S. <i>et al.</i> (2021) Overexpression of the ubiquitin-editing enzyme A20 in the brain
lesions of Multiple Sclerosis patients: moving from systemic to central nervous system
inflammation. Brain Pathol. 31 (2): 283-96.
This product is shipped at ambient temperature. It is recommended to aliquot and store at
-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for
short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
Avoid repeated freezing and thawing as this may denature the antibody. Storage in
frost-free freezers is not recommended.
Guaranteed until date of expiry. Please see product label.

Health And Safety Material Safety Datasheet documentation #10053 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA1815 10053

Storage

Guarantee

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>			
Goat Anti Mouse IgG (STAR76)	RPE		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>		
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,		
	DyLight®650, DyLight®680, DyLight®800,		
	DyLight®650, DyLight®680, DyLight®800, FITC, HRP		
Rabbit Anti Mouse IgG (STAR9)			
Rabbit Anti Mouse IgG (STAR9) Goat Anti Mouse IgG (STAR77)	FITC, HRP		
o ()	FITC, HRP FITC		

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M418605:230427'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint