

Datasheet: MCA1815T

BATCH NUMBER 151806

Description:	MOUSE ANTI HUMAN CD68
Specificity:	CD68
Other names:	MACROSIALIN
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	514H12
Isotype:	IgG2a
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/40 - 1/80
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting		▪		
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Tris/EDTA buffer pH 9.0 is recommended for this purpose. *N.B. The epitope recognised by clone 514H12 is sensitive to peroxide. It is therefore recommended that any peroxidase-blocking steps be performed after incubation with this antibody.*

Target Species	Human
Product Form	Tissue Culture Supernatant - liquid

Preservative Stabilisers	0.09% Sodium Azide
Immunogen	Fusion protein corresponding to external domain of human CD68.
External Database Links	<p>UniProt: P34810 Related reagents</p> <p>Entrez Gene: 968 CD68 Related reagents</p>
RRID	AB_2074721
Fusion Partners	Spleen cells from immunized mice were fused with cells of the mouse p3-NS1-Ag4-1 myeloma cell line
Specificity	Mouse anti Human CD68 antibody, clone 514H12 recognizes the human CD68 cell surface antigen, a ~110 kDa glycoprotein primarily expressed by macrophages and monocytes.
Histology Positive Control Tissue	Human Tonsil
References	<ol style="list-style-type: none"> Hameed, A. <i>et al.</i> (1994) Immunohistochemical expression of CD68 antigen in human peripheral blood T cells. Hum Pathol. 25 (9): 872-6. Madden, L.R. <i>et al.</i> (2010) Proangiogenic scaffolds as functional templates for cardiac tissue engineering. Proc Natl Acad Sci U S A. 107 (34): 15211-6. Muthana, M. <i>et al.</i> (2011) Use of macrophages to target therapeutic adenovirus to human prostate tumors. Cancer Res. 71 (5): 1805-15. da Costa, C.E. <i>et al.</i> (2005) Presence of osteoclast-like multinucleated giant cells in the bone and nonostotic lesions of Langerhans cell histiocytosis. J Exp Med. 201 (5): 687-93. Wang, X. <i>et al.</i> (2006) Monocyte/macrophage and T-cell infiltrates in peritoneum of patients with ovarian cancer or benign pelvic disease. J Transl Med. 4: 30. Coury, F. <i>et al.</i> (2008) Langerhans cell histiocytosis reveals a new IL-17A-dependent pathway of dendritic cell fusion. Nat Med. 14: 81-7. Rodriguez-Agudo, D. <i>et al.</i> (2006) Localization of StarD5 cholesterol binding protein. J Lipid Res. 47: 1168-75. Achard, V. <i>et al.</i> (2007) Renin receptor expression in human adipose tissue. Am J Physiol Regul Integr Comp Physiol. 292: R274-82. Paulmyer-Lacroix, O. <i>et al.</i> (2006) Expression of adrenomedullin in adipose tissue of lean and obese women. Eur J Endocrinol. 155: 177-85. Hever, A. <i>et al.</i> (2007) Human endometriosis is associated with plasma cells and overexpression of B lymphocyte stimulator. Proc Natl Acad Sci U S A. 104: 12451-6. Silaghi, A. <i>et al.</i> (2007) Expression of adrenomedullin in human epicardial adipose tissue: role of coronary status. Am J Physiol Endocrinol Metab. 293: E1443-50. Kaibara, N. <i>et al.</i> (2008) Comparative histopathological analysis between tenosynovitis and joint synovitis in rheumatoid arthritis. Histopathology. 52: 856-64. Moskovszky, L. <i>et al.</i> (2009) Genomic instability in giant cell tumor of bone. A study of

52 cases using DNA ploidy, relocalization FISH, and array-CGH analysis. [Genes Chromosomes Cancer. 48: 468-79.](#)

14. Angel, C.E. *et al.* (2007) CD14+ antigen-presenting cells in human dermis are less mature than their CD1a+ counterparts. [Int Immunol. 19: 1271-9.](#)

15. de Vos van Steenwijk PJ *et al.* (2013) Tumor-infiltrating CD14-positive myeloid cells and CD8-positive T-cells prolong survival in patients with cervical carcinoma. [Int J Cancer. 133 \(12\): 2884-94.](#)

16. Quispel, W.T. *et al.* (2016) Tertiary lymphoid structures are confined to patients presenting with unifocal Langerhans Cell Histiocytosis. [Oncoimmunology. 5 \(8\): e1164364.](#)

17. Toh, M.L. *et al.* (2014) Bone- and cartilage-protective effects of a monoclonal antibody against colony-stimulating factor 1 receptor in experimental arthritis. [Arthritis Rheumatol. 66 \(11\): 2989-3000.](#)

18. Schirmer, L. *et al.* (2014) Differential loss of KIR4.1 immunoreactivity in multiple sclerosis lesions. [Ann Neurol. 75 \(6\): 810-28.](#)

19. Schrevel, M. *et al.* (2017) Autocrine expression of the epidermal growth factor receptor ligand heparin-binding EGF-like growth factor in cervical cancer. [Int J Oncol. 50 \(6\): 1947-54.](#)

20. Zhang, W. *et al.* (2013) Myeloid clusters are associated with a pro-metastatic environment and poor prognosis in smoking-related early stage non-small cell lung cancer. [PLoS One. 8 \(5\): e65121.](#)

21. Bobosha, K. *et al.* (2014) T-cell regulation in lepromatous leprosy. [PLoS Negl Trop Dis. 8 \(4\): e2773.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

Guaranteed until date of expiry. Please see product label.

Health And Safety Information

Material Safety Datasheet documentation #10055 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1815T10055>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M353473:190423'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)