

Datasheet: MCA1821F BATCH NUMBER 152974

Description:	HAMSTER ANTI MOUSE CD79b:FITC
Specificity:	CD79b
Other names:	B29
Format:	FITC
Product Type:	Monoclonal Antibody
Product Type: Clone:	Monoclonal Antibody HM79-11
	•

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Dete	ermined	Suggested Dilution	
	Flow Cytometry				Neat	
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/postive controls.					
Target Species	Mouse					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max (nn	n) Emission I	Max (nm)		
	FITC	490	52			
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin					
Approx. Protein	IgG concentration 0.1m	ıg/ml				

Concentrations

immunogen CD79 alpha/CD79 beta heterodimer purified from WEHI-231 B cells. External Database Links UniProt: P15530 Related reagents Synonyms Igb RND AB_323287 Fusion Partners Lymph node cells from immunised Armenian hamsters were fused with cells of the mouse X63-Ag8 653 myeloma cell line. Specificity Hamster anti Mouse CD79b antibody, clone HM79-11 recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ^d cells in 100ul. The Fc region of monocional antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBicoc FCR (BUF04TA/B). References 1.Koyama, M. et al. (1997) CD79 alpha/CD79 beta heterodiment CD79 ao nimmature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10); er76115. 3. Jellnek, F. et al. (2005) Characteristics of two mouse bor-abi-transformed cell lines. II. Pathological lesions induced in mice. Folla Biol (Praha), 51 (4); 93-102. 4. Ankeny, D. P. et al. (2005) Characteristics of two mouse bor-abi-transformed cell lines. II. Pathological lesions induc				
Links UniProt: P15530 Related reagents Synonyms Igb RRID AB_323287 Fusion Partners Lymph node cells from immunised Armenian hamsters were fused with cells of the mouse X63-Ag8.653 myeloma cell line. Specificity Hamster anti Mouse CD79b antibody, clone HM79-11 recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a Coll specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc receptors. This may be reduced by using SeroBlock. FcR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/D279 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9(11): 1767-72. 1. Logan, D. <i>et al.</i> (2005) Characteristics of two mouse bor-abi-transformed cell lines. II. Pathological lesions induced in mice. Folla Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. <i>et al.</i> (2005) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem, 99 (4): 1073-87. 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the	Immunogen	CD79 alpha/CD79 beta heterodimer purified from WEHI-231 B cells.		
RRID AB_323287 Fusion Partners Lymph node cells from immunised Armenian hamsters were fused with cells of the mouse X63-Ag8.653 myeloma cell line. Specificity Hamster anti Mouse CD79b antibody, clone HM79-11 recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FCR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9(11): 1767-72. 2. Luger, D. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 39: 102. 4. Ankeny, D. P. <i>et al.</i> (2005) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B (mphocyte activation and lupus-like autoantibody synthesis. <u>J Neurochem. 99 (4): 1073-87.</u> 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 16		P15530 Related reagents Entrez Gene:		
Fusion Partners Lymph node cells from immunised Armenian hamsters were fused with cells of the mouse X63-Ag8.653 myeloma cell line. Specificity Hamster anti Mouse CD79b antibody, clone HM79-11 recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUE041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72, 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. <u>PLoS One. 8 (10): e76115.</u> 3. Jellnek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 903-102. 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem. 99 (4): 1073-87. 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a	Synonyms	lgb		
X63-Ag8.653 myeloma cell line. Specificity Hamster anti Mouse CD79b antibody, clone HM79-11 recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B). References 1. Koyama, M. et al. (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72. 2. Luger, D. et al. (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): e76115. 3. Jellnek, F. et al. (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. et al. (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem, 99 (4): 1073-87. 5. Lacoste-collin, L. et al. (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymph	RRID	AB_323287		
expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer). Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72, 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): e76115, 3. Jellnek, F, <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102, 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem, 99 (4): 1073-87, 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.	Fusion Partners			
immunoglobulin, and is expressed throughout B cell differentiation. CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72. 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): <u>e76115.</u> 3. Jelínek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem. 99 (4): 1073-87. 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.	Specificity	expressed by B cells as part of the B cell receptor complex (immunoglobulin and the		
stages. Flow Cytometry Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B). References 1. Koyama, M. et al. (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72. 2. Luger, D. et al. (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): e76115. 3. Jelínek, F. et al. (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. et al. (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem. 99 (4): 1073-87. 5. Lacoste-collin, L. et al. (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.				
The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72. 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): e76115. 3. Jelínek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem. 99 (4): 1073-87. 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.				
affinity Fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/B</u>). References 1. Koyama, M. <i>et al.</i> (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. Int Immunol. 9 (11): 1767-72. 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. PLoS One. 8 (10): e76115. 3. Jelínek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102. 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. J Neurochem. 99 (4): 1073-87. 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.	Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.		
 cell surfaces without associated mu heavy chain. <u>Int Immunol. 9 (11): 1767-72.</u> 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. <u>PLoS One. 8 (10): e76115.</u> 3. Jelínek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. <u>Folia Biol (Praha). 51 (4): 93-102.</u> 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. <u>J Neurochem. 99 (4): 1073-87.</u> 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. <u>Radiat Res. 168 (6): 725-32.</u> 				
Storage Store at +4°C or at -20°C if preferred.	References	 cell surfaces without associated mu heavy chain. <u>Int Immunol. 9 (11): 1767-72.</u> 2. Luger, D. <i>et al.</i> (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. <u>PLoS One. 8 (10): e76115.</u> 3. Jelínek, F. <i>et al.</i> (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. <u>Folia Biol (Praha). 51 (4): 93-102.</u> 4. Ankeny, D.P. <i>et al.</i> (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. <u>J Neurochem. 99 (4): 1073-87.</u> 5. Lacoste-collin, L. <i>et al.</i> (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. 		
	Storage			

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1821F 10041	
Regulatory	For research purposes only	

Related Products

Recommended Negative Controls

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:FITC (MCA2356F)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	id.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M365754:200529'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint