

Datasheet: MCA1821F

Description:	HAMSTER ANTI MOUSE CD79b:FITC		
Specificity:	CD79b		
Other names:	B29		
Format:	FITC		
<b>Product Type:</b>	Monoclonal Antibody		
Clone:	HM79-11		
Isotype:	IgG		
Quantity:	0.1 mg		

# **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

lm	m	un	Oa	en
		u	vы	<b>CII</b>

CD79 alpha/CD79 beta heterodimer purified from WEHI-231 B cells.

# External Database Links

#### **UniProt**:

P15530 Related reagents

#### **Entrez Gene:**

15985 Cd79b Related reagents

#### **Synonyms**

lgb

#### RRID

AB 323287

### **Fusion Partners**

Lymph node cells from immunized Armenian hamsters were fused with cells of the mouse X63-Ag8.653 myeloma cell line.

### **Specificity**

**Hamster anti Mouse CD79b antibody, clone HM79-11** recognizes murine CD79 beta, expressed by B cells as part of the B cell receptor complex (immunoglobulin and the CD79 alpha/beta heterodimer).

Murine CD79 beta is expressed at the cell surface of pro-B cells prior to surface immunoglobulin, and is expressed throughout B cell differentiation.

CD79 beta is a B cell specific marker, valuable for the detection of B cells at all maturation stages.

### Flow Cytometry

Use  $10\mu l$  of the suggested working dilution to label  $10^6$  cells in  $100\mu l$ .

The Fc region of monoclonal antibodies may bind to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/BUF041B</u>).

### References

- 1. Koyama, M. *et al.* (1997) CD79 alpha/CD79 beta heterodimers are expressed on pro-B cell surfaces without associated mu heavy chain. <u>Int Immunol. 9 (11): 1767-72.</u>
- 2. Jelínek, F. *et al.* (2005) Characteristics of two mouse bcr-abl-transformed cell lines. II. Pathological lesions induced in mice. Folia Biol (Praha). 51 (4): 93-102.
- 3. Ankeny, D.P. *et al.* (2006) Spinal cord injury triggers systemic autoimmunity: evidence for chronic B lymphocyte activation and lupus-like autoantibody synthesis. <u>J Neurochem.</u> 99 (4): 1073-87.
- 4. Lacoste-collin, L. *et al.* (2007) Effect of continuous irradiation with a very low dose of gamma rays on life span and the immune system in SJL mice prone to B-cell lymphoma. Radiat Res. 168 (6): 725-32.
- 5. Luger, D. *et al.* (2013) Expression of the B-cell receptor component CD79a on immature myeloid cells contributes to their tumor promoting effects. <u>PLoS One. 8 (10):</u> <u>e76115.</u>

### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1821F">https://www.bio-rad-antibodies.com/SDS/MCA1821F</a> 10041
Regulatory	For research purposes only

# **Related Products**

# **Recommended Negative Controls**

# HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:FITC (MCA2356F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M411664:221104'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint