

Datasheet: MCA1824PET

Description:	MOUSE ANTI HUMAN CD89:RPE
Specificity:	CD89
Other names:	Fc ALPHA R
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MIP8a
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute in 0.25 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Immunogen	Recombinant soluble human Fc alpha R.		
External Database Links	UniProt: P24071 Related reagents		
	Entrez Gene: 2204 FCAR Related reagents		

Synonyms	CD89
Specificity	<p>Mouse anti Human CD89 antibody, clone MIP8a recognizes the human CD89 cell surface antigen, a 50-75 kDa cell surface glycoprotein that is also known as the IgA receptor (Fc alpha R).</p> <p>CD89 is expressed by peripheral blood monocytes and neutrophils.</p> <p>MIP8a blocks binding of IgA to the Fc alpha R, and also inhibits neutrophil phagocytosis of IgA complexes.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	<ol style="list-style-type: none"> 1. Zhang, W. <i>et al.</i> (2000) Neutrophil lactoferrin release induced by IgA immune complexes differed from that induced by cross-linking of fcalpha receptors (FcalphaR) with a monoclonal antibody, MIP8a. Clin Exp Immunol. 121 (1): 106-11. 2. Lu, J. <i>et al.</i> (2011) Recognition and functional activation of the human IgA receptor (Fc{alpha}RI) by C-reactive protein. Proc Natl Acad Sci U S A. 108: 4974-9. 3. Duc, M. <i>et al.</i> (2010) Antigen binding to secretory immunoglobulin A results in decreased sensitivity to intestinal proteases and increased binding to cellular Fc receptors. J Biol Chem. 285: 953-60. 4. Wu, J. <i>et al.</i> (2007) FcαRI (CD89) alleles determine the proinflammatory potential of serum IgA. J Immunol. 178: 3973-82. 5. Hamre, R. <i>et al.</i> (2003) Expression and modulation of the human immunoglobulin A Fc receptor (CD89) and the FcR gamma chain on myeloid cells in blood and tissue. Scand J Immunol. 57: 506-16. 6. Qian, K. <i>et al.</i> (2008) Functional expression of IgA receptor FcalphaRI on human platelets. J Leukoc Biol. 84: 1492-500. 7. Van Egmond, M. (2011) Method for the treatment or prophylaxis of chronic inflammatory diseases. European Patent Application No: 12/736963 8. Pascal, V. <i>et al.</i> (2012) Anti-CD20 IgA can protect mice against lymphoma development: evaluation of the direct impact of IgA and cytotoxic effector recruitment on CD20 target cells. Haematologica. 97 (11): 1686-94. 9. Mladenov, R. <i>et al.</i> (2015) The Fc-alpha receptor is a new target antigen for immunotherapy of myeloid leukemia. Int J Cancer. 137 (11): 2729-38. 10. Aleyd, E. <i>et al.</i> (2016) IgA Complexes in Plasma and Synovial Fluid of Patients with Rheumatoid Arthritis Induce Neutrophil Extracellular Traps via FcαRI. J Immunol. 197 (12): 4552-9.
Storage	<p>Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A)

HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M323049:180727'

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)