

Datasheet: MCA1847FT

BATCH NUMBER 1710

Description:	MOUSE ANTI HUMAN CD81:FITC	
Specificity:	CD81	
Other names:	TAPA-1	
Format:	FITC	
Product Type:	Monoclonal Antibody	
Clone:	1D6	
Isotype:	lgG1	
Quantity:	25 μg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Species Cross
Reactivity

Target Species

Reacts with: Chimpanzee, Sheep, Goat

Human

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Dr	~~	uct	E	rm
Pr	oa	uct	FO	rm

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	OCI-LY8 cells aggregated by 5A6 (another CD81 antibody)
External Database	UniProt:
Links	P60033 Related reagents
	Entrez Gene: 975 CD81 Related reagents
	<u>373</u> OD01 <u>Related reagents</u>
Synonyms	TAPA1, TSPAN28
RRID	AB_1102391
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse PX3-Ag.8.653 myeloma cell line
Specificity	Mouse anti Human CD81 antibody, clone 1D6 recognizes human CD81, a 26 kDa cell surface antigen also known as TAPA-1, and a member of the tetraspanin family. CD81 is widely expressed on human leucocytes and appears to be involved in a variety of cellular leucocytes including activation, proliferation and differentiation.
	Mouse anti Human CD81 antibody, clone 1D6 is a potent CD81 reagent, induces homotypic adhesion and has powerful anti-proliferative effects.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Schick, M.R. & Levy, S. (1993) The TAPA-1 molecule is associated on the surface of B cells with HLA-DR molecules. J Immunol. 151 (8): 4090-7. Levy, S. et al. (1998) CD81 (TAPA-1): a molecule involved in signal transduction and cell adhesion in the immune system. Annu Rev Immunol. 16: 89-109. Griebel, P.J. et al. (2007) Cross-reactivity of mAbs to human CD antigens with sheep leukocytes. Vet Immunol Immunopathol. 119: 115-22. Welton, J.L. et al (2010) Proteomics analysis of bladder cancer exosomes. Mol Cell Proteomics. 9: 1324-38. Davis, W.C. et al. (2007) Use of flow cytometry to identify monoclonal antibodies that recognize conserved epitopes on orthologous leukocyte differentiation antigens in goats, llamas, and rabbits. Vet Immunol Immunopathol. 119: 123-30. Flint, M. et al. (1999) Characterization of hepatitis C virus E2 glycoprotein interaction with a putative cellular receptor, CD81. J Virol. 73:6235-44.
	7. Parthasarathy, V. et al. (2009) Distinct roles for tetraspanins CD9, CD63 and CD81 in

the formation of multinucleated giant cells. Immunology. 127: 237-48.

- 8. Rohlena, J. et al. (2009) Endothelial CD81 is a marker of early human atherosclerotic plaques and facilitates monocyte adhesion. Cardiovasc Res. 81: 187-96.
- 9. Ventress, J.K. et al. (2016) Peptides from Tetraspanin CD9 Are Potent Inhibitors of Staphylococcus Aureus Adherence to Keratinocytes. PLoS One. 11 (7): e0160387.
- 10. Mleczko, J. et al. (2018) Extracellular Vesicles from Hypoxic Adipocytes and Obese Subjects Reduce Insulin-Stimulated Glucose Uptake. Mol Nutr Food Res. 62 (5)Feb 20 [Epub ahead of print].

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1847FT 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365789:200529'

Printed on 19 Mar 2025