

Datasheet: MCA1848

BATCH NUMBER 167502

Description:	MOUSE ANTI HUMAN CD11a
Specificity:	CD11a
Other names:	INTEGRIN ALPHA L CHAIN, LFA-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	38
Isotype:	IgG2a
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% sodium azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1 mg/ml
External Database Links	<p>UniProt: P20701 Related reagents</p> <p>Entrez Gene: 3683 ITGAL Related reagents</p>
Synonyms	CD11A
RRID	AB_322231
Specificity	<p>Mouse anti Human CD11a antibody, clone 38 recognizes the integrin alpha L subunit, also known as CD11a, leukocyte function-associated molecule 1 alpha chain LFA-1 or integrin alpha-L. CD11a is a ~180 kDa glycoprotein expressed by all human leucocytes. CD11a is involved in cell-cell adhesion processes and binds to ICAM-1, 2 and 3.</p> <p>Mouse anti Human CD11a antibody, clone 38 inhibits CD11a mediated adhesion of T cells to ICAM-1 (Stewart <i>et al.</i> 1996).</p>
Histology Positive Control Tissue	Human tonsil
References	<ol style="list-style-type: none"> 1. Dransfield, I. & Hogg, N. (1989) Regulated expression of Mg2+ binding epitope on leukocyte integrin alpha subunits. EMBO J. 8 (12): 3759-65. 2. Berg, L.P. <i>et al.</i> (2002) Functional consequences of noncognate interactions between CD4+ memory T lymphocytes and the endothelium. J Immunol. 168 (7): 3227-34. 3. Foster, G.R. <i>et al.</i> (2004) IFN-alpha subtypes differentially affect human T cell motility. J Immunol. 173: 1663-70. 4. Evans, B.J. <i>et al.</i> (2006) Shedding of lymphocyte function-associated antigen-1 (LFA-1) in a human inflammatory response. Blood. 107: 3593-9. 5. Suchard, S.J. <i>et al.</i> (2010) An LFA-1 (alphaLbeta2) small-molecule antagonist reduces inflammation and joint destruction in murine models of arthritis. J Immunol. 184: 3917-26. 6. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. FASEB J. 23: 492-502. 7. Daehn, I.S. (2010) T-lymphocyte-induced, fas-mediated apoptosis is associated with early keratinocyte differentiation. Exp Dermatol. 19: 372-80. 8. Dominical, V.M. <i>et al.</i> (2014) Prominent role of platelets in the formation of circulating neutrophil-red cell heterocellular aggregates in sickle cell anemia. Haematologica. 99 (11): e214-7. 9. Stax, A.M. <i>et al.</i> (2017) Autoreactivity to Sulfatide by Human Invariant NKT Cells. J Immunol. 199 (1): 97-106.

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1848 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M411801:221107'

Printed on 18 Jan 2024