

Datasheet: MCA1848GA

BATCH NUMBER 169289

Description:	MOUSE ANTI HUMAN CD11a
Specificity:	CD11a
Other names:	INTEGRIN ALPHA L CHAIN, LFA-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	38
Isotype:	lgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/50 - 1/100
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation				
Western Blotting				
Immunofluorescence				
Functional Assays				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% sodium azide (NaN ₃)	
Carrier Free	Yes	
Approx. Protein Concentrations	IgG concentration 1 mg/ml	
External Database Links	UniProt: P20701 Related reagents	
	Entrez Gene: 3683 ITGAL Related reagents	
Synonyms	CD11A	
RRID	AB_324104	
Specificity	Mouse anti Human CD11a antibody, clone 38 recognizes the integrin alpha L subunit, also known as CD11a, leukocyte function-associated molecule 1 alpha chain LFA-1 or integrin alpha-L. CD11a is a ~180 kDa glycoprotein expressed by all human leucocytes. CD11a is involved in cell-cell adhesion processes and binds to ICAM-1, 2 and 3.	
	Mouse anti Human CD11a antibody, clone 38 inhibits CD11a mediated adhesion of T cells to ICAM-1 (Stewart <i>et al.</i> 1996).	
Histology Positive Control Tissue	Human tonsil	
References	1. Dransfield, I. & Hogg, N. (1989) Regulated expression of Mg2+ binding epitope on leukocyte integrin alpha subunits. EMBO J. 8 (12): 3759-65. 2. Berg, L.P. <i>et al.</i> (2002) Functional consequences of noncognate interactions between CD4+ memory T lymphocytes and the endothelium. J Immunol. 168 (7): 3227-34. 3. Foster, G.R. <i>et al.</i> (2004) IFN-alpha subtypes differentially affect human T cell motility. J Immunol. 173: 1663-70. 4. Evans, B.J. <i>et al.</i> (2006) Shedding of lymphocyte function-associated antigen-1 (LFA-1) in a human inflammatory response. Blood. 107: 3593-9 . 5. Suchard, S.J. <i>et al.</i> (2010) An LFA-1 (alphaLbeta2) small-molecule antagonist reduces inflammation and joint destruction in murine models of arthritis. J Immunol. 184: 3917-26 . 6. Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. FASEB J. 23: 492-502 . 7. Daehn, I.S. (2010) T-lymphocyte-induced, fas-mediated apoptosis is associated with early keratinocyte differentiation. Exp Dermatol. 19: 372-80 . 8. Dominical, V.M. <i>et al.</i> (2014) Prominent role of platelets in the formation of circulating neutrophil-red cell heterocellular aggregates in sickle cell anemia. Haematologica. 99 (11): e214-7 . 9. Stax, A.M. <i>et al.</i> (2017) Autoreactivity to Sulfatide by Human Invariant NKT Cells. J Immunol. 199 (1): 97-106 .	

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1848GA 10040		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) **RPE** Goat Anti Mouse IgG IgA IgM (STAR87...) HRP Goat Anti Mouse IgG (STAR76...) **RPE** Goat Anti Mouse IgG (STAR70...) **FITC**

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) **HRP**

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) **FITC** Goat Anti Mouse IgG (STAR77...) **HRP**

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 America

Fax: +44 (0)1865 852 739 Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com

Worldwide

Tel: +49 (0) 89 8090 95 21 Europe Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M432555:240903'

Tel: +44 (0)1865 852 700

Printed on 03 Sep 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint