

Datasheet: MCA1848XZ

Description:	MOUSE ANTI HUMAN CD11a: Preservative Free
Specificity:	CD11a
Other names:	INTEGRIN ALPHA L CHAIN, LFA-1
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	38
Isotype:	lgG2a
Quantity:	1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry				1/50 - 1/100		
	Immunohistology - Frozen						
	Immunohistology - Paraffin			•			
	ELISA			•			
	Immunoprecipitation	-					
	Western Blotting			•			
	Immunofluorescence						
	Functional Assays						
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.						
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant			n tissue culture			
Buffer Solution	Phosphate buffered salin	e					
Preservative Stabilisers	None present						

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: P20701 Related reagents Entrez Gene: <u>3683</u> ITGAL <u>Related reagents</u>
Synonyms	CD11A
RRID	AB_322927
Specificity	Mouse anti Human CD11a antibody, clone 38 recognizes the integrin alpha L subunit, also known as CD11a, leukocyte function-associated molecule 1 alpha chain LFA-1 or integrin alpha-L. CD11a is a ~180 kDa glycoprotein expressed by all human leucocytes. CD11a is involved in cell-cell adhesion processes and binds to ICAM-1, 2 and 3. Mouse anti Human CD11a antibody, clone 38 inhibits CD11a mediated adhesion of T cells
	to ICAM-1 (Stewart <i>et al.</i> 1996).
References	 Dransfield, I. & Hogg, N. (1989) Regulated expression of Mg2+ binding epitope on leukocyte integrin alpha subunits. <u>EMBO J. 8 (12): 3759-65.</u> Berg, L.P. <i>et al.</i> (2002) Functional consequences of noncognate interactions between CD4+ memory T lymphocytes and the endothelium. <u>J Immunol. 168 (7): 3227-34.</u> Foster, G.R. <i>et al.</i> (2004) IFN-alpha subtypes differentially affect human T cell motility. <u>J Immunol. 173: 1663-70.</u> Evans, B.J. <i>et al.</i> (2006) Shedding of lymphocyte function-associated antigen-1 (LFA-1) in a human inflammatory response. <u>Blood. 107: 3593-9.</u> Suchard, S.J. <i>et al.</i> (2010) An LFA-1 (alphaLbeta2) small-molecule antagonist reduces inflammation and joint destruction in murine models of arthritis. <u>J Immunol. 184: 3917-26.</u> Porter, J.C. and Hall, A. (2009) Epithelial ICAM-1 and ICAM-2 regulate the egression of human T cells across the bronchial epithelium. <u>FASEB J. 23: 492-502.</u> Daehn, I.S. (2010) T-lymphocyte-induced, fas-mediated apoptosis is associated with early keratinocyte differentiation. <u>Exp Dermatol. 19: 372-80.</u> Dominical, V.M. <i>et al.</i> (2014) Prominent role of platelets in the formation of circulating neutrophil-red cell heterocellular aggregates in sickle cell anemia. <u>Haematologica. 99 (11): e214-7.</u> Stax, A.M. <i>et al.</i> (2017) Autoreactivity to Sulfatide by Human Invariant NKT Cells. <u>J Immunol. 199 (1): 97-106.</u>
Storage	Store at -20°C only. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we

	recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10162 available at: https://www.bio-rad-antibodies.com/SDS/MCA1848XZ 10162
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Ar	nti Mouse IgG (STAR12)	RPE					
Goat Ant	i Mouse IgG (STAR70)	<u>FITC</u>					
Goat Ant	Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>						
Rabbit Ar	nti Mouse IgG (STAR9)	<u>FITC</u>					
Goat Ant	i Mouse IgG (STAR76)	RPE					
Goat Ant	i Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLig DyLight®650, DyLight®680, Dy					
		FITC, HRP					
Rabbit Ar	nti Mouse IgG (STAR13)	HRP					
Goat Anti Mouse IgG (Fc) (STAR120) <u>FITC</u> , <u>HRP</u>							
Recommended Negative Controls							
MOUSE IgG2a NEGATIVE CONTROL (MCA929)							
North & South America	Tel: +1 800 265 7376 Worldwi Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	de Tel: +44 (0)1865 852 700 E Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.co	- Fax	: +49 (0) 89 8090 95 21 x: +49 (0) 89 8090 95 50 nail: antibody_sales_de@bio-rad.com			
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M411800:221107'							

Printed on 03 Sep 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint