

Datasheet: MCA1849

BATCH NUMBER 148285

Description:	RAT ANTI MOUSE MARCO
Specificity:	MARCO
Other names:	SCAVENGER RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	ED31
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry		▪		
Immunohistology - Frozen	▪			1/200 - 1/400
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen CHO cells expressing mouse MARCO.

External Database Links

UniProt:

[Q60754](#) [Related reagents](#)

Entrez Gene:

[17167](#) Marco [Related reagents](#)

RRID AB_322923

Fusion Partners Spleen cells from immunised rats were fused with cells of the mouse SP2/0 myeloma cell line.

Specificity

Rat anti Mouse MARCO antibody, clone ED31 recognizes the murine cell surface antigen designated MARCO (Macrophage receptor with collagenous structure), which is a member of the class A scavenger receptor family.

MARCO is expressed by distinct populations of macrophages in the spleen and lymph nodes, but is rapidly induced on macrophages in other tissues (e.g. Liver Kupffer cells) during infection or LPS treatment.

Rat anti Mouse MARCO antibody, clone ED31 binds to the C-terminal cysteine rich domain of MARCO, and has been shown to block ligand binding.

References

1. van der Laan, L.J. *et al.* (1999) Regulation and functional involvement of macrophage scavenger receptor MARCO in clearance of bacteria *in vivo*. [J Immunol. 162 \(2\): 939-47.](#)
2. van der Laan, L.J. *et al.* (1997) Macrophage scavenger receptor MARCO: *in vitro* and *in vivo* regulation and involvement in the anti-bacterial host defense. [Immunol Lett. 57 \(1-3\): 203-8.](#)
3. Dioszeghy, V. *et al.* (2008) 12/15-Lipoxygenase regulates the inflammatory response to bacterial products *in vivo*. [J Immunol. 181: 6514-24.](#)
4. Devey, L. *et al.* (2009) Tissue-resident macrophages protect the liver from ischemia reperfusion injury via a heme oxygenase-1-dependent mechanism. [Mol Ther. 17: 65-72.](#)
5. Hsu, K.M. *et al.* (2009) Murine cytomegalovirus displays selective infection of cells within hours after systemic administration. [J Gen Virol. 90: 33-43.](#)
6. Kang, Y.S. *et al.* (2004) The C-type lectin SIGN-R1 mediates uptake of the capsular polysaccharide of *Streptococcus pneumoniae* in the marginal zone of mouse spleen. [Proc Natl Acad Sci U S A. 101: 215-20.](#)
7. Karlsson, M.C. *et al.* (2003) Macrophages control the retention and trafficking of B lymphocytes in the splenic marginal zone. [J Exp Med. 198: 333-40.](#)
8. Taylor, P.R. *et al.* (2004) Development of a specific system for targeting protein to

- metallophilic macrophages. [Proc Natl Acad Sci U S A. 101: 1963-8.](#)
9. Alarcón, R. *et al.* (2005) Expression of scavenger receptors in glial cells. Comparing the adhesion of astrocytes and microglia from neonatal rats to surface-bound beta-amyloid. [J Biol Chem. 280: 30406-15.](#)
10. Anthony, R.M. *et al.* (2008) Identification of a receptor required for the anti-inflammatory activity of IVIG. [Proc Natl Acad Sci U S A. 105: 19571-8.](#)
11. Beamer, C.A. and Holian. A. (2008) Silica suppresses Toll-like receptor ligand-induced dendritic cell activation. [FASEB J. 22: 2053-63.](#)
12. Granucci, F. *et al.* (2003) The scavenger receptor MARCO mediates cytoskeleton rearrangements in dendritic cells and microglia. [Blood. 102: 2940-7.](#)
13. Mattsson, J. *et al.* (2011) Complement activation and complement receptors on follicular dendritic cells are critical for the function of a targeted adjuvant. [J Immunol. 187: 3641-52.](#)
14. Fukui Y *et al.* (2013) Effect of *Lactobacillus brevis* KB290 on the cell-mediated cytotoxic activity of mouse splenocytes: a DNA microarray analysis. [Br J Nutr. 110 \(9\): 1617-29.](#)
15. Hayashi, M. *et al.* (2017) Advax, a Delta Inulin Microparticle, Potentiates In-built Adjuvant Property of Co-administered Vaccines. [EBioMedicine. 15: 127-36.](#)
16. Martinez, N. *et al.* (2016) Impaired Recognition of *Mycobacterium tuberculosis* by Alveolar Macrophages From Diabetic Mice. [J Infect Dis. 214 \(11\): 1629-1637.](#)
17. Parsa, R. *et al.* (2016) BAFF-secreting neutrophils drive plasma cell responses during emergency granulopoiesis. [J Exp Med. 213 \(8\): 1537-53.](#)
18. Marrella, V. *et al.* (2015) IL-10 critically modulates B cell responsiveness in Rankl-/- mice. [J Immunol. 194 \(9\): 4144-53.](#)
19. Kolan, S.S. *et al.* (2015) Lack of non-hematopoietic SIRPα signaling disturbs the splenic marginal zone architecture resulting in accumulation and displacement of marginal zone B cells. [Biochem Biophys Res Commun. 460 \(3\): 645-50.](#)
20. Flores, M> *et al.* (2015) FcγRIIB prevents inflammatory type I IFN production from plasmacytoid dendritic cells during a viral memory response. [J Immunol. 194 \(9\): 4240-50.](#)
21. Zhao, Z. *et al.* (2019) Neutrophil-Derived MRP14 Supports Plasma Cell Commitment and Protects Myeloma Cells from Apoptosis. [J Immunol Res. 2019: 9561350.](#)
22. Hamasaki, M. *et al.* (2020) Transcriptional profiling of murine macrophages stimulated with cartilage fragments revealed a strategy for treatment of progressive osteoarthritis. [Sci Rep. 10 \(1\): 7558.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1849>
10040

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: antibody_sales_us@bio-rad.com**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M365798:200529'

Printed on 31 May 2024