

## Datasheet: MCA1853F

**BATCH NUMBER 163733**

<b>Description:</b>	MOUSE ANTI HUMAN CD163:FITC
<b>Specificity:</b>	CD163
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	EDHu-1
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Human

**Species Cross Reactivity** Reacts with: Rhesus Monkey, Sheep, Pig, Guinea Pig, Bovine, Cynomolgus monkey  
**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

**Product Form** Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.1mg/ml
<b>Immunogen</b>	Leucocytes harvested from the pleural cavity of patients with idiopathic spontaneous pneumothorax
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q86VB7</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">9332</a>    CD163    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	M130
<b>RRID</b>	AB_2260020
<b>Specificity</b>	<p><b>Mouse anti Human CD163 antibody, clone EDHu-1</b> recognizes the human CD163 cell surface antigen, a 130-140 kDa glycoprotein expressed by tissue macrophages. CD163 expression may be induced on monocytes by culture in dexamethasone.</p> <p>Clone EDHu-1 is reported to inhibit the binding of haptoglobin/hemoglobin to CD163 (<a href="#">Madsen <i>et al.</i> 2004</a>). Truncation mutation analysis demonstrates binding of EDHu-1 occurs via the N-terminal region of CD163 containing the first three scavenger receptor, Cysteine-rich, <a href="#">SRCR domains</a> the third domain being critical as, cleavage of this domain at the major cleavage site <a href="#">ASP-265</a> abrogates binding to the N-terminal fragment.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Kristiansen, M. <i>et al.</i> (2001) Identification of the haemoglobin scavenger receptor. <a href="#">Nature. 409 (6817): 198-201.</a></li> <li>Madsen, M. <i>et al.</i> (2004) Molecular characterization of the haptoglobin.hemoglobin receptor CD163. Ligand binding properties of the scavenger receptor cysteine-rich domain region. <a href="#">J Biol Chem. 279 (49): 51561-7.</a></li> <li>Kim, W.K. <i>et al.</i> (2006) CD163 identifies perivascular macrophages in normal and viral encephalitic brains and potential precursors to perivascular macrophages in blood. <a href="#">Am J Pathol. 168 (3): 822-34.</a></li> <li>Moreno JA <i>et al.</i> (2010) Peripheral artery disease is associated with a high CD163/TWEAK plasma ratio. <a href="#">Arterioscler Thromb Vasc Biol. 30 (6): 1253-62.</a></li> <li>Herrmann-Hoesing, L.M. (2010) Ovine progressive pneumonia virus capsid antigen as found in CD163- and CD172a-positive alveolar macrophages of persistently infected sheep. <a href="#">Vet Pathol. 47: 518-28.</a></li> <li>Asleh, R. <i>et al.</i> (2003) Genetically determined heterogeneity in hemoglobin scavenging and susceptibility to diabetic cardiovascular disease. <a href="#">Circ Res. 92: 1193-200.</a></li> <li>Fabriek, B.O. <i>et al.</i> (2007) The macrophage CD163 surface glycoprotein is an</li> </ol>

- erythroblast adhesion receptor. [Blood 109: 5223-9.](#)
8. Jensen, T.O. *et al.* (2009) Macrophage markers in serum and tumor have prognostic impact in American Joint Committee on Cancer stage I/II melanoma. [J Clin Oncol. 27: 3330-7.](#)
  9. Montes de Oca, M. *et al.* (2005) Skeletal muscle inflammation and nitric oxide in patients with COPD. [Eur Respir J. 26: 390-7.](#)
  10. Martens JH *et al.* (2006) Differential expression of a gene signature for scavenger/lectin receptors by endothelial cells and macrophages in human lymph node sinuses, the primary sites of regional metastasis. [J Pathol. 208 \(4\): 574-89.](#)
  11. Vinet-Oliphant, H. *et al.* (2010) Neurokinin-1 receptor (NK1-R) expression in the brains of SIV-infected rhesus macaques: implications for substance P in NK1-R immune cell trafficking into the CNS. [Am J Pathol. 177: 1286-97.](#)
  12. Wang, X. *et al.* (2006) Monocyte/macrophage and T-cell infiltrates in peritoneum of patients with ovarian cancer or benign pelvic disease. [J Transl Med. 4: 30.](#)
  13. Grund, S. *et al.* (2009) The microglial/macrophagic response at the tumour-brain border of invasive meningiomas. [Neuropathol Appl Neurobiol. 35: 82-8.](#)
  14. Jorgensen, J.M. *et al.* (2009) Expression level, tissue distribution pattern, and prognostic impact of vascular endothelial growth factors VEGF and VEGF-C and their receptors Flt-1, KDR, and Flt-4 in different subtypes of non-Hodgkin lymphomas. [Leuk Lymphoma. 50: 1647-60.](#)
  15. Moreno, J.A. *et al.* (2009) The CD163-expressing macrophages recognize and internalize TWEAK: potential consequences in atherosclerosis. [Atherosclerosis. 207: 103-10.](#)
  16. Tang, Z. *et al.* (2013) Glucocorticoids Enhance CD163 Expression in Placental Hofbauer Cells. [Endocrinology 154: 471-82.](#)
  17. Boyle, J.J. *et al.* (2009) Coronary intraplaque hemorrhage evokes a novel atheroprotective macrophage phenotype. [Am J Pathol. 174: 1097-108.](#)
  18. Taus, N.S. *et al.* (2010) Sheep (*Ovis aries*) airway epithelial cells support ovine herpesvirus 2 lytic replication in vivo. [Vet Microbiol. 145: 47-53.](#)
  19. Seeboth, J. *et al.* (2012) The fungal T-2 toxin alters the activation of primary macrophages induced by TLR-agonists resulting in a decrease of the inflammatory response in the pig. [Vet Res. 43: 35.](#)
  20. Berglin, L. *et al.* (2014) *In situ* characterization of intrahepatic non-parenchymal cells in PSC reveals phenotypic patterns associated with disease severity. [PLoS One 9: e105375.](#)
  21. Liu, J. *et al.* (2014) Evidence for mTOR pathway activation in a spectrum of epilepsy-associated pathologies. [Acta Neuropathol Commun. 2: 71.](#)
  22. Baek, J.H. *et al.* (2014) Extracellular Hb enhances cardiac toxicity in endotoxemic guinea pigs: protective role of haptoglobin. [Toxins \(Basel\) 6: 1244-59.](#)
  23. Micci, L. *et al.* (2014) CD4 depletion in SIV-infected macaques results in macrophage and microglia infection with rapid turnover of infected cells. [PLoS Pathog. 10: e1004467.](#)
  24. Piriã E *et al.* (2015) Macrophages modulate migration and invasion of human tongue squamous cell carcinoma. [PLoS One 10 \(3\): e0120895.](#)
  25. Arranz-Solís D *et al.* (2016) Systemic and local immune responses in sheep after *Neospora caninum* experimental infection at early, mid and late gestation. [Vet Res. 47 \(1\): 2.](#)
  26. Lakritz, J R. *et al.* (2016)  $\alpha$ 4-Integrin Antibody Treatment Blocks

- Monocyte/Macrophage Traffic to, Vascular Cell Adhesion Molecule-1 Expression in, and Pathology of the Dorsal Root Ganglia in an SIV Macaque Model of HIV-Peripheral Neuropathy. [Am J Pathol. 186 \(7\): 1754-61.](#)
27. Fry, L.M. *et al.* (2016) East Coast Fever Caused by *Theileria parva* Is Characterized by Macrophage Activation Associated with Vasculitis and Respiratory Failure. [PLoS One 11 \(5\): e0156004.](#)
28. Schultz, N. *et al.* (2017) Amylin alters human brain pericyte viability and NG2 expression. [J Cereb Blood Flow Metab. 37 \(4\): 1470-82.](#)
29. Zhang, W. *et al.* (2013) Myeloid clusters are associated with a pro-metastatic environment and poor prognosis in smoking-related early stage non-small cell lung cancer. [PLoS One 8: e65121.](#)
30. Furukawa S *et al.* (2017) Interleukin-33 produced by M2 macrophages and other immune cells contributes to Th2 immune reaction of IgG4-related disease. [Sci Rep. 7: 42413.](#)
31. Farina, A. *et al.* (2017) Epstein-Barr virus lytic infection promotes activation of Toll-like receptor 8 innate immune response in systemic sclerosis monocytes. [Arthritis Res Ther. 19 \(1\): 39.](#)
32. Mallard, J. *et al.* (2017) A method for obtaining simian immunodeficiency virus RNA sequences from laser capture microdissected and immune captured CD68+ and CD163+ macrophages from frozen tissue sections of bone marrow and brain. [J Immunol Methods. 442: 59-63.](#)
33. Blair, T.C. *et al.* (2016) Immunopathology of Japanese macaque encephalomyelitis is similar to multiple sclerosis. [J Neuroimmunol. 291: 1-10.](#)
34. Zhu, C. *et al.* (2017) Activation of CECR1 in M2-like TAMs promotes paracrine stimulation-mediated glial tumor progression. [Neuro Oncol. 19 \(5\): 648-59.](#)
35. Derricott, H. *et al.* (2016) Characterizing Villitis of Unknown Etiology and Inflammation in Stillbirth. [Am J Pathol. 186 \(4\): 952-61.](#)
36. Wächter, C. *et al.* (2016) Loss of cerebellar neurons in the progression of lentiviral disease: effects of CNS-permeant antiretroviral therapy. [J Neuroinflammation. 13 \(1\): 272.](#)
37. Chen, J. *et al.* (2019) Generation of Pigs Resistant to Highly Pathogenic-Porcine Reproductive and Respiratory Syndrome Virus through Gene Editing of CD163. [Int J Biol Sci. 15 \(2\): 481-492.](#)
38. Palaiologou, E. *et al.* (2020) Human placental villi contain stromal macrovesicles associated with networks of stellate cells. [J Anat. 236 \(1\): 132-41.](#)
39. Kong, L.Q. *et al.* (2013) The clinical significance of the CD163+ and CD68+ macrophages in patients with hepatocellular carcinoma. [PLoS One. 8 \(3\): e59771.](#)
40. Zhao, S. *et al.* (2020) CD14<sup>+</sup> monocytes and CD163<sup>+</sup> macrophages correlate with the severity of liver fibrosis in patients with chronic hepatitis C [Experimental and Therapeutic Medicine. 20 \(6\): 1-1.](#)
41. Hayashi, K. *et al.* (2020) The Natural History of Spontaneously Occurred Endometriosis in Cynomolgus Monkeys by Monthly Follow-Up Laparoscopy for Two Years. [Tohoku J Exp Med. 251 \(4\): 241-53.](#)
42. Gonçalves, V.M. *et al.* (2021) Macrophage and Lymphocyte Infiltration Is Associated with Volumetric Tumor Size but Not with Volumetric Growth in the T&yym;l;bingen Schwannoma Cohort. [Cancers \(Basel\). 13 \(3\): 466.](#)
43. Pestronk, A. (2020) Chronic Graft Versus Host Myopathies: Noninflammatory, Multi-Tissue Pathology With Glycosylation Disorders. [J Neuropathol Exp Neurol. 79 \(1\): 102-12.](#)

44. Holsapple, J.S. *et al.* (2021) Low Intensity Shockwave Treatment Modulates Macrophage Functions Beneficial to Healing Chronic Wounds. [Int J Mol Sci. 22\(15\):7844.](#)
45. da Silva M.C.M. *et al.* (2021) Inhibition of CSF1R, a receptor involved in microglia viability, alters behavioral and molecular changes induced by cocaine. [Sci Rep. 11 \(1\): 15989.](#)
46. Frafjord, A. *et al.* (2020) Antibody combinations for optimized staining of macrophages in human lung tumours. [Scand J Immunol. 92 \(1\): e12889.](#)
47. Ma, H. *et al.* (2021) Structural comparison of CD163 SRCR5 from different species sheds some light on its involvement in porcine reproductive and respiratory syndrome virus-2 infection *in vitro.* [Vet Res. 52 \(1\): 97.](#)
48. Kemmerer, C.L. *et al.* (2021) Cerebrospinal fluid cytokine levels are associated with macrophage infiltration into tumor tissues of glioma patients [BMC Cancer. 21\(1\):1108.](#)
49. Lehmann, M. *et al.* (2021) New onset of mainly guttate psoriasis after COVID-19 vaccination: a case report. [J Eur Acad Dermatol Venereol. 35 \(11\): e752-e755.](#)
50. Baumann, D. *et al.* (2021) p38 MAPK signaling in M1 macrophages results in selective elimination of M2 macrophages by MEK inhibition. [J Immunother Cancer.9 \(7\): e002319.](#)
51. Ikezumi, Y. *et al.* (2021) Steroid treatment promotes an M2 anti-inflammatory macrophage phenotype in childhood lupus nephritis. [Pediatr Nephrol. 36 \(2\): 349-59.](#)
52. Eligini, S. *et al.* (2019) Biological profile of monocyte-derived macrophages in coronary heart disease patients: implications for plaque morphology. [Sci Rep. 9 \(1\): 8680.](#)
53. Edwards, J.H. *et al.* (2021) Integration and functional performance of a decellularised porcine superflexor tendon graft in an ovine model of anterior cruciate ligament reconstruction. [Biomaterials. 279: 121204.](#)
54. Ströbel, S. *et al.* (2021) A 3D primary human cell-based in vitro model of non-alcoholic steatohepatitis for efficacy testing of clinical drug candidates. [Sci Rep. 11 \(1\): 22765.](#)
55. Galea, I. *et al.* (2022) Iron Deposition in the Brain After Aneurysmal Subarachnoid Hemorrhage. [Stroke. 53 \(5\): 1633-42.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1853F>  
10041

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

## Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M385233:210513'

Printed on 11 Apr 2024

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)