

Datasheet: MCA1853F BATCH NUMBER 163733

Description:	MOUSE ANTI HUMAN CD163:FITC
Specificity:	CD163
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	EDHu-1
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	information. For genera	•				
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat - 1/10	
	Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross Reactivity	Reacts with: Rhesus Monkey, Sheep, Pig, Guinea Pig, Bovine, Cynomolgus monkey N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.					
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)		
	FITC	490		525		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant					
Buffer Solution	Phosphate buffered sa	lline				

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1mg/ml
Immunogen	Leucocytes harvested from the pleural cavity of patients with idiopathic spontaneous pneumothorax
External Database Links	UniProt: Q86VB7 Related reagents Entrez Gene: 9332 CD163 Related reagents
Synonyms	M130
RRID	AB_2260020
Specificity	 Mouse anti Human CD163 antibody, clone EDHu-1 recognizes the human CD163 cell surface antigen, a 130-140 kDa glycoprotein expressed by tissue macrophages. CD163 expression may be induced on monocytes by culture in dexamethasone. Clone EDHu-1 is reported to inhibit the binding of haptoglobin/hemoglobin to CD163 (Madsen <i>et al.</i> 2004).Truncation mutation analysis demonstrates binding of EDHu-1 occurs via the N-terminal region of CD163 containing the first three scavenger receptor, Cysteine-rich, <u>SRCR domains</u> the third domain being critical as, cleavage of this domain at the major cleavage site <u>ASP-265</u> abrogates binding to the N-terminal fragment.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Kristiansen, M. <i>et al.</i> (2001) Identification of the haemoglobin scavenger receptor. Nature. 409 (6817): 198-201. Madsen, M. <i>et al.</i> (2004) Molecular characterization of the haptoglobin.hemoglobin receptor CD163. Ligand binding properties of the scavenger receptor cysteine-rich domain region. J Biol Chem. 279 (49): 51561-7. Kim, W.K. <i>et al.</i> (2006) CD163 identifies perivascular macrophages in normal and viral encephalitic brains and potential precursors to perivascular macrophages in blood. Am J Pathol. 168 (3): 822-34. Moreno JA <i>et al.</i> (2010) Peripheral artery disease is associated with a high CD163/TWEAK plasma ratio. Arterioscler Thromb Vasc Biol. 30 (6): 1253-62. Herrmann-Hoesing, L.M. (2010) Ovine progressive pneumonia virus capsid antigen as found in CD163- and CD172a-positive alveolar macrophages of persistently infected sheep. Vet Pathol. 47: 518-28. Asleh, R. <i>et al.</i> (2003) Genetically determined heterogeneity in hemoglobin scavenging and susceptibility to diabetic cardiovascular disease. <u>Circ Res. 92: 1193-200.</u> Fabriek, B.O. <i>et al.</i> (2007) The macrophage CD163 surface glycoprotein is an

erythroblast adhesion receptor. Blood 109: 5223-9.

8. Jensen, T.O. *et al.* (2009) Macrophage markers in serum and tumor have prognostic impact in American Joint Committee on Cancer stage I/II melanoma. <u>J Clin Oncol. 27:</u> <u>3330-7.</u>

9. Montes de Oca, M. *et al.* (2005) Skeletal muscle inflammation and nitric oxide in patients with COPD. <u>Eur Respir J. 26: 390-7.</u>

10. Martens JH *et al.* (2006) Differential expression of a gene signature for scavenger/lectin receptors by endothelial cells and macrophages in human lymph node sinuses, the primary sites of regional metastasis. <u>J Pathol. 208 (4): 574-89.</u>

11. Vinet-Oliphant, H. *et al.* (2010) Neurokinin-1 receptor (NK1-R) expression in the brains of SIV-infected rhesus macaques: implications for substance P in NK1-R immune cell trafficking into the CNS. <u>Am J Pathol. 177: 1286-97.</u>

12. Wang, X. *et al.* (2006) Monocyte/macrophage and T-cell infiltrates in peritoneum of patients with ovarian cancer or benign pelvic disease. <u>J Transl Med. 4: 30.</u>

13. Grund, S. *et al* (2009) The microglial/macrophagic response at the tumour-brain border of invasive meningiomas. <u>Neuropathol Appl Neurobiol. 35: 82-8.</u>

14. Jorgensen, J.M. *et al* (2009) Expression level, tissue distribution pattern, and prognostic impact of vascular endothelial growth factors VEGF and VEGF-C and their receptors Flt-1, KDR, and Flt-4 in different subtypes of non-Hodgkin lymphomas. <u>Leuk Lymphoma. 50: 1647-60.</u>

15. Moreno, J.A. *et al.* (2009) The CD163-expressing macrophages recognize and internalize TWEAK: potential consequences in atherosclerosis. <u>Atherosclerosis. 207:</u> 103-10.

16. Tang, Z. *et al.* (2013) Glucocorticoids Enhance CD163 Expression in Placental Hofbauer Cells. <u>Endocrinology 154: 471-82.</u>

17. Boyle, J.J. *et al.* (2009) Coronary intraplaque hemorrhage evokes a novel atheroprotective macrophage phenotype. <u>Am J Pathol. 174: 1097-108.</u>

18. Taus, N.S. *et al.* (2010) Sheep (Ovis aries) airway epithelial cells support ovine herpesvirus 2 lytic replication in vivo. <u>Vet Microbiol. 145: 47-53.</u>

19. Seeboth, J. *et al.* (2012) The fungal T-2 toxin alters the activation of primary macrophages induced by TLR-agonists resulting in a decrease of the inflammatory response in the pig. <u>Vet Res. 43: 35.</u>

20. Berglin, L. *et al.* (2014) *In situ* characterization of intrahepatic non-parenchymal cells in PSC reveals phenotypic patterns associated with disease severity. <u>PLoS One 9:</u> <u>e105375.</u>

21. Liu, J. *et al.* (2014) Evidence for mTOR pathway activation in a spectrum of epilepsyassociated pathologies. <u>Acta Neuropathol Commun. 2: 71.</u>

22. Baek, J.H. *et al.* (2014) Extracellular Hb enhances cardiac toxicity in endotoxemic guinea pigs: protective role of haptoglobin. <u>Toxins (Basel) 6: 1244-59.</u>

23. Micci, L, *et al.* (2014) CD4 depletion in SIV-infected macaques results in macrophage and microglia infection with rapid turnover of infected cells. <u>PLoS Pathog. 10: e1004467</u>.
24. Pirilä E *et al.* (2015) Macrophages modulate migration and invasion of human tongue squamous cell carcinoma. PLoS One 10 (3): e0120895.

25. Arranz-Solís D *et al.* (2016) Systemic and local immune responses in sheep after *Neospora caninum* experimental infection at early, mid and late gestation. <u>Vet Res. 47 (1):</u> 2.

26. Lakritz, J R. et al. (2016) α4-Integrin Antibody Treatment Blocks

Monocyte/Macrophage Traffic to, Vascular Cell Adhesion Molecule-1 Expression in, and Pathology of the Dorsal Root Ganglia in an SIV Macaque Model of HIV-Peripheral Neuropathy. <u>Am J Pathol. 186 (7): 1754-61.</u>

27. Fry, L.M. *et al.* (2016) East Coast Fever Caused by *Theileria parva* Is Characterized by Macrophage Activation Associated with Vasculitis and Respiratory Failure. <u>PLoS One</u> <u>11 (5): e0156004.</u>

28. Schultz, N. *et al.* (2017) Amylin alters human brain pericyte viability and NG2 expression. <u>J Cereb Blood Flow Metab. 37 (4): 1470-82.</u>

29. Zhang, W. *et al.* (2013) Myeloid clusters are associated with a pro-metastatic environment and poor prognosis in smoking-related early stage non-small cell lung cancer. <u>PLoS One 8: e65121.</u>

30. Furukawa S *et al.* (2017) Interleukin-33 produced by M2 macrophages and other immune cells contributes to Th2 immune reaction of IgG4-related disease. <u>Sci Rep. 7:</u> <u>42413.</u>

31. Farina, A. *et al.* (2017) Epstein-Barr virus lytic infection promotes activation of Toll-like receptor 8 innate immune response in systemic sclerosis monocytes. <u>Arthritis Res Ther.</u> <u>19 (1): 39.</u>

32. Mallard, J. *et al.* (2017) A method for obtaining simian immunodeficiency virus RNA sequences from laser capture microdissected and immune captured CD68+ and CD163+ macrophages from frozen tissue sections of bone marrow and brain. <u>J Immunol Methods.</u> <u>442: 59-63.</u>

33. Blair, T.C. *et al.* (2016) Immunopathology of Japanese macaque encephalomyelitis is similar to multiple sclerosis. <u>J Neuroimmunol. 291: 1-10.</u>

34. Zhu, C. *et al.* (2017) Activation of CECR1 in M2-like TAMs promotes paracrine stimulation-mediated glial tumor progression. <u>Neuro Oncol. 19 (5): 648-59.</u>

35. Derricott, H. *et al.* (2016) Characterizing Villitis of Unknown Etiology and Inflammation in Stillbirth. <u>Am J Pathol. 186 (4): 952-61.</u>

36. Wächter, C. *et al.* (2016) Loss of cerebellar neurons in the progression of lentiviral disease: effects of CNS-permeant antiretroviral therapy. <u>J Neuroinflammation</u>. 13 (1): 272.
37. Chen, J. *et al.* (2019) Generation of Pigs Resistant to Highly Pathogenic-Porcine Reproductive and Respiratory Syndrome Virus through Gene Editing of CD163. <u>Int J Biol</u> Sci. 15 (2): 481-492.

38. Palaiologou, E. *et al.* (2020) Human placental villi contain stromal macrovesicles associated with networks of stellate cells. <u>J Anat. 236 (1): 132-41.</u>

39. Kong, L.Q. *et al.* (2013) The clinical significance of the CD163+ and CD68+ macrophages in patients with hepatocellular carcinoma. <u>PLoS One. 8 (3): e59771.</u>

40. Zhao, S. *et al.* (2020) CD14⁺ monocytes and CD163⁺ macrophages correlate with the severity of liver fibrosis in patients with chronic hepatitis C <u>Experimental and Therapeutic</u> <u>Medicine. 20 (6): 1-1.</u>

41. Hayashi, K. *et al.* (2020) The Natural History of Spontaneously Occurred Endometriosis in Cynomolgus Monkeys by Monthly Follow-Up Laparoscopy for Two Years. <u>Tohoku J Exp Med. 251 (4): 241-53.</u>

42. Gonçalves, V.M. *et al.* (2021) Macrophage and Lymphocyte Infiltration Is Associated with Volumetric Tumor Size but Not with Volumetric Growth in the T&yyml;bingen Schwannoma Cohort. <u>Cancers (Basel). 13 (3): 466.</u>

43. Pestronk, A. (2020) Chronic Graft Versus Host Myopathies: Noninflammatory, Multi-Tissue Pathology With Glycosylation Disorders. <u>J Neuropathol Exp Neurol. 79 (1): 102-12.</u>

	 44. Holsapple, J.S. <i>et al.</i> (2021) Low Intensity Shockwave Treat Macrophage Functions Beneficial to Healing Chronic Wounds. J. 45. da Silva M.C.M. <i>et al.</i> (2021) Inhibition of CSF1R, a receptor viability, alters behavioral and molecular changes induced by contability, alters behavioral and molecular changes in the state space. State alters, and the contability of the contabilit	Int J Mol Sci. 22(15):7844. or involved in microglia ocaine. Sci Rep. 11 (1): ed staining of macrophages 5 from different species 4 respiratory vels are associated with <u>IC Cancer. 21(1):1108.</u> riasis after COVID-19 <u>11): e752-e755.</u> ophages results in selective <u>er Cancer.9 (7): e002319.</u> anti-inflammatory <u>ephrol. 36 (2): 349-59.</u> d macrophages in coronary <u>Rep. 9 (1): 8680.</u> mance of a decellularised ruciate ligament
Storage	This product is shipped at ambient temperature. It is recommen -20°C on receipt. When thawed, aliquot the sample as needed. short term use (up to 4 weeks) and store the remaining aliquots	Keep aliquots at 2-8°C for at -20°C.
	Avoid repeated freezing and thawing as this may denature the a frost-free freezers is not recommended. This product is photose protected from light.	, ,
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1853F 10041	
Regulatory	For research purposes only	

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@b	bio-rad.com	Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385233:210513'

Printed on 29 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint