

Datasheet: MCA1855

Description:	MOUSE ANTI HUMAN CD161
Specificity:	CD161
Other names:	NKR-P1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	B199.2
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting		▪		

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified human NK cells cultured in IL-2 (Bennett et al. 1996)
External Database Links	<p>UniProt: Q12918 Related reagents</p> <p>Entrez Gene: 3820 KLRB1 Related reagents</p>
Synonyms	CLEC5B, NKRP1A
RRID	AB_323331
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse P2X63.Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human CD161 antibody, clone B199.2 recognizes the human Killer cell lectin-like receptor subfamily B member 1, also known as CD161, C-type lectin domain family 5 member B, HNKRP-1a, NKRP-1A or Natural killer cell surface protein P1A. CD161 is a 225 amino acid ~25 kDa predicted molecular mass, single pass type II transmembrane glycoprotein with a single C-type lectin domain. CD161 is expressed by almost all NK cells and a subset of CD3+ve T cells (Lanier 1994).</p> <p>CD161, a member of the C-lectin is expressed as a disulphide bond-linked homodimeric cell surface protein, comprising two chains of ~40-44 kDa (Lanier et al. 1994). CD161 acts as a receptor for another c-type lectin, LLT1 with roles in the regulation of NK cell and T cell function (Aldemir et al. 2005).</p> <p>Mouse anti Human CD161 antibody, clone B199.2 cross-competes with and recognizes a similar epitope to the DX1 monoclonal antibody (Lanier et al. 1994).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Bennett, I.M. <i>et al.</i> (1996) Definition of a natural killer NKR-P1A+/CD56-/CD16- functionally immature human NK cell subset that differentiates <i>in vitro</i> in the presence of interleukin 12. J Exp Med. 184 (5): 1845-56. Azzoni, L. <i>et al.</i> (1998) Differential transcriptional regulation of CD161 and a novel gene, 197/15a, by IL-2, IL-15, and IL-12 in NK and T cells. J Immunol. 161 (7): 3493-500. de Lalla, C. <i>et al.</i> (2011) Invariant NKT Cell Reconstitution in Pediatric Leukemia Patients Given HLA-Haploidentical Stem Cell Transplantation Defines Distinct CD4+ and CD4- Subset Dynamics and Correlates with Remission State. J Immunol. 186: 4490-9. Huarte, E. <i>et al.</i> (2008) PILAR is a novel modulator of human T-cell expansion. Blood. 112: 1259-68.

5. Williams, P.J. *et al.* (2009) Altered decidual leucocyte populations in the placental bed in pre-eclampsia and foetal growth restriction: a comparison with late normal pregnancy. [Reproduction. 138: 177-84.](#)
6. Higai, K. *et al.* (2006) Binding of sialyl Lewis X antigen to lectin-like receptors on NK cells induces cytotoxicity and tyrosine phosphorylation of a 17-kDa protein. [Biochim Biophys Acta. 1760 \(9\): 1355-63.](#)
7. Birchall, M.A. *et al.* (2008) Immunologic response of the laryngeal mucosa to extraesophageal reflux. [Ann Otol Rhinol Laryngol. 117: 891-5.](#)
8. Bossard, C. *et al.* (2012) Plasmacytoid dendritic cells and Th17 immune response contribution in gastrointestinal acute graft-versus-host disease. [Leukemia. 26: 1471-4.](#)
9. Richter, J. *et al.* (2010) CD161 receptor participates in both impairing NK cell cytotoxicity and the response to glycans and vimentin in patients with rheumatoid arthritis. [Clin Immunol. 136: 139-47.](#)
10. Abrahamsson, S.V. *et al.* (2013) Non-myeloablative autologous haematopoietic stem cell transplantation expands regulatory cells and depletes IL-17 producing mucosal-associated invariant T cells in multiple sclerosis. [Brain. 136: 2888-903.](#)
11. Rother, S. *et al.* (2015) The c.503T>C Polymorphism in the Human KLRB1 Gene Alters Ligand Binding and Inhibitory Potential of CD161 Molecules. [PLoS One. 10 \(8\): e0135682.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M411860:221108'

Printed on 25 Jan 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)