

Datasheet: MCA1855PET

Description:	MOUSE ANTI HUMAN CD161:RPE
Specificity:	CD161
Other names:	NKR-P1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	B199.2
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute in 0.25 ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Immunogen	Purified human NK cells cultured in IL-2 (Bennett et al. 1996)		
External Database Links	UniProt:		
	Q12918	Related reagents	

Entrez Gene:

[3820](#) KLRB1 [Related reagents](#)

Synonyms CLEC5B, NKR P1A

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse P2X63.Ag8.653 myeloma cell line.

Specificity **Mouse anti Human CD161 antibody, clone B199.2** recognizes the human Killer cell lectin-like receptor subfamily B member 1, also known as CD161, C-type lectin domain family 5 member B, HNKR-P1a, NKR-P1A or Natural killer cell surface protein P1A. CD161 is a 225 amino acid ~25 kDa predicted molecular mass, single pass type II transmembrane glycoprotein with a single [C-type lectin](#) domain. CD161 is expressed by almost all NK cells and a subset of CD3+ve T cells ([Lanier 1994](#)).

CD161, a member of the C-lectin is expressed as a disulphide bond-linked homodimeric cell surface protein, comprising two chains of ~40-44 kDa ([Lanier et al. 1994](#)). CD161 acts as a receptor for another c-type lectin, LLT1 with roles in the regulation of NK cell and T cell function ([Aldemir et al. 2005](#)).

Mouse anti Human CD161 antibody, clone B199.2 cross-competes with and recognizes a similar epitope to the DX1 monoclonal antibody ([Lanier et al. 1994](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Bennett, I.M. *et al.* (1996) Definition of a natural killer NKR-P1A+/CD56-/CD16- functionally immature human NK cell subset that differentiates *in vitro* in the presence of interleukin 12. [J Exp Med. 184 \(5\): 1845-56.](#)
2. Azzoni, L. *et al.* (1998) Differential transcriptional regulation of CD161 and a novel gene, 197/15a, by IL-2, IL-15, and IL-12 in NK and T cells. [J Immunol. 161 \(7\): 3493-500.](#)
3. de Lalla, C. *et al.* (2011) Invariant NKT Cell Reconstitution in Pediatric Leukemia Patients Given HLA-Haploidentical Stem Cell Transplantation Defines Distinct CD4+ and CD4- Subset Dynamics and Correlates with Remission State. [J Immunol. 186: 4490-9.](#)
4. Huarte, E. *et al.* (2008) PILAR is a novel modulator of human T-cell expansion. [Blood. 112: 1259-68.](#)
5. Williams, P.J. *et al.* (2009) Altered decidual leucocyte populations in the placental bed in pre-eclampsia and foetal growth restriction: a comparison with late normal pregnancy. [Reproduction. 138: 177-84.](#)
6. Higai, K. *et al.* (2006) Binding of sialyl Lewis X antigen to lectin-like receptors on NK cells induces cytotoxicity and tyrosine phosphorylation of a 17-kDa protein. [Biochim Biophys Acta. 1760: 1355-63.](#)
7. Birchall, M.A. *et al.* (2008) Immunologic response of the laryngeal mucosa to extraesophageal reflux. [Ann Otol Rhinol Laryngol. 117: 891-5.](#)
8. Bossard, C. *et al.* (2012) Plasmacytoid dendritic cells and Th17 immune response contribution in gastrointestinal acute graft-versus-host disease. [Leukemia. 26: 1471-4.](#)
9. Higai, K. *et al.* (2008) Binding of sialyl Lewis X antigen to lectin-like receptors on NK cells induces cytotoxicity and tyrosine phosphorylation of a 17-kDa protein [Biochim Biophys Acta. 1760: 1355-63.](#)
10. Richter, J. *et al.* (2010) CD161 receptor participates in both impairing NK cell cytotoxicity and the response to glycans and vimentin in patients with rheumatoid arthritis. [Clin Immunol. 136: 139-47.](#)
11. Abrahamsson, S.V. *et al.* (2013) Non-myeloablative autologous haematopoietic stem cell

transplantation expands regulatory cells and depletes IL-17 producing mucosal-associated invariant T cells in multiple sclerosis. [Brain. 136: 2888-903.](#)

12. Rother S *et al.* (2015) The c.503T>C Polymorphism in the Human KLRB1 Gene Alters Ligand Binding and Inhibitory Potential of CD161 Molecules. [PLoS One. 10 \(8\): e0135682.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

12 months from date of reconstitution.

Health And Safety Information

Material Safety Datasheet documentation #10075 available at: 10075: <https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M323085:180727'

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)