

# Datasheet: MCA1870FT

**BATCH NUMBER 151044**

<b>Description:</b>	MOUSE ANTI HUMAN CD326:FITC
<b>Specificity:</b>	CD326
<b>Other names:</b>	Ep-CAM
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	VU-1D9
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN <sub>3</sub> )		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	0.1mg/ml		

<b>Immunogen</b>	HG9 cell line (small cell lung carcinoma)
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P16422</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">4072</a>    EPCAM    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TROP1
<b>RRID</b>	AB_1101938
<b>Fusion Partners</b>	Spleen cells from immunised mice were fused with cells of the mouse SP2/0 myeloma cell line
<b>Specificity</b>	<p><b>Mouse anti Human CD326 antibody, clone VU-1D9</b> recognizes epithelial cell adhesion molecule (Ep-CAM), a ~34 kDa cell surface antigen otherwise known as CD326, MOC31 or Ber-EP4. CD326 is a type 1 transmembrane glycoprotein, expressed on the basolateral cell membrane of the majority of epithelial tissues, with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells.</p> <p>Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells, and <i>de novo</i> expression being seen in dysplastic squamous epithelium</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Koga, Y. <i>et al.</i> (2008) Improved recovery of exfoliated colonocytes from feces using newly developed immunomagnetic beads. <a href="#">Gastroenterol Res Pract. 2008: 605273.</a></li> <li>2. Fillmore, C.M. <i>et al.</i> (2010) Estrogen expands breast cancer stem-like cells through paracrine FGF/Tbx3 signaling. <a href="#">Proc Natl Acad Sci U S A. 107 (50): 21737-42.</a></li> <li>3. Murakata, A. <i>et al.</i> (2011) Gene expression signature of the gross morphology in hepatocellular carcinoma. <a href="#">Ann Surg. 253: 94-100.</a></li> <li>4. Keller, P.J. <i>et al.</i> (2012) Defining the cellular precursors to human breast cancer. <a href="#">Proc Natl Acad Sci U S A. 109 (8): 2772-7.</a></li> <li>5. Jeong, H.T. <i>et al.</i> (2012) MRI features of hepatocellular carcinoma expressing progenitor cell markers. <a href="#">Liver Int. 32: 430-40.</a></li> <li>6. Armstrong, A.J. <i>et al.</i> (2019) Pharmacodynamic study of radium-223 in men with bone metastatic castration resistant prostate cancer. <a href="#">PLoS One. 14 (5): e0216934.</a></li> </ol>
<b>Further Reading</b>	1. Winter, M.J. <i>et al.</i> (2003) The epithelial cell adhesion molecule (Ep-CAM) as a morphoregulatory molecule is a tool in surgical pathology. <a href="#">Am J Pathol. 163 (6): 2139-48.</a>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p>

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1870FT">https://www.bio-rad-antibodies.com/SDS/MCA1870FT</a> 10041
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M365839:200529'

**Printed on 18 Jan 2024**