

Datasheet: MCA1870FT

BATCH NUMBER 1711

Description:	MOUSE ANTI HUMAN CD326:FITC
Specificity:	CD326
Other names:	Ep-CAM
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	VU-1D9
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)		
	1% Bovine Serum Albumin		
Approx. Protein Concentrations	0.1mg/ml		

Immunogen	HG9 cell line (small cell lung carcinoma)
External Database Links	<p>UniProt: P16422 Related reagents</p> <p>Entrez Gene: 4072 EPCAM Related reagents</p>
Synonyms	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TROP1
RRID	AB_1101938
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse SP2/0 myeloma cell line
Specificity	<p>Mouse anti Human CD326 antibody, clone VU-1D9 recognizes epithelial cell adhesion molecule (Ep-CAM), a ~34 kDa cell surface antigen otherwise known as CD326, MOC31 or Ber-EP4. CD326 is a type 1 transmembrane glycoprotein, expressed on the basolateral cell membrane of the majority of epithelial tissues, with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells.</p> <p>Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells, and <i>de novo</i> expression being seen in dysplastic squamous epithelium</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> 1. Koga, Y. <i>et al.</i> (2008) Improved recovery of exfoliated colonocytes from feces using newly developed immunomagnetic beads. Gastroenterol Res Pract. 2008: 605273. 2. Fillmore, C.M. <i>et al.</i> (2010) Estrogen expands breast cancer stem-like cells through paracrine FGF/Tbx3 signaling. Proc Natl Acad Sci U S A. 107 (50): 21737-42. 3. Murakata, A. <i>et al.</i> (2011) Gene expression signature of the gross morphology in hepatocellular carcinoma. Ann Surg. 253: 94-100. 4. Keller, P.J. <i>et al.</i> (2012) Defining the cellular precursors to human breast cancer. Proc Natl Acad Sci U S A. 109 (8): 2772-7. 5. Jeong, H.T. <i>et al.</i> (2012) MRI features of hepatocellular carcinoma expressing progenitor cell markers. Liver Int. 32: 430-40. 6. Armstrong, A.J. <i>et al.</i> (2019) Pharmacodynamic study of radium-223 in men with bone metastatic castration resistant prostate cancer. PLoS One. 14 (5): e0216934.
Further Reading	1. Winter, M.J. <i>et al.</i> (2003) The epithelial cell adhesion molecule (Ep-CAM) as a morphoregulatory molecule is a tool in surgical pathology. Am J Pathol. 163 (6): 2139-48.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p>

This product should be stored undiluted. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1870FT 10041
--------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M365839:200529'

Printed on 18 Jan 2024