

Datasheet: MCA1870GT

BATCH NUMBER 151424

Description:	MOUSE ANTI HUMAN CD326
Specificity:	CD326
Other names:	Ep-CAM
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	VU-1D9
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/200
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	1 mg/ml
Immunogen	HG9 cell line (small cell lung carcinoma)
External Database Links	<p>UniProt: P16422 Related reagents</p> <p>Entrez Gene: 4072 EPCAM Related reagents</p>
Synonyms	GA733-2, M1S2, M4S1, MIC18, TACSTD1, TROP1
RRID	AB_2098800
Fusion Partners	Spleen cells from immunised mice were fused with cells of the mouse SP2/0 myeloma cell line
Specificity	<p>Mouse anti Human CD326 antibody, clone VU-1D9 recognizes epithelial cell adhesion molecule (Ep-CAM), a ~34 kDa cell surface antigen otherwise known as CD326, MOC31 or Ber-EP4. CD326 is a type 1 transmembrane glycoprotein, expressed on the basolateral cell membrane of the majority of epithelial tissues, with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells.</p> <p>Ep-CAM expression has been reported to be a possible marker of early malignancy, with expression being increased in tumor cells, and <i>de novo</i> expression being seen in dysplastic squamous epithelium</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
Histology Positive Control Tissue	Large Bowel
Western Blotting	MCA1870G detects a band of approximately 38kDa in HT-29 cell lysates
References	<ol style="list-style-type: none"> 1. Koga, Y. <i>et al.</i> (2008) Improved recovery of exfoliated colonocytes from feces using newly developed immunomagnetic beads. Gastroenterol Res Pract. 2008: 605273. 2. Fillmore, C.M. <i>et al.</i> (2010) Estrogen expands breast cancer stem-like cells through paracrine FGF/Tbx3 signaling. Proc Natl Acad Sci U S A. 107 (50): 21737-42. 3. Murakata, A. <i>et al.</i> (2011) Gene expression signature of the gross morphology in hepatocellular carcinoma. Ann Surg. 253: 94-100.

4. Keller, P.J. *et al.* (2012) Defining the cellular precursors to human breast cancer. [Proc Natl Acad Sci U S A. 109 \(8\): 2772-7.](#)
5. Jeong, H.T. *et al.* (2012) MRI features of hepatocellular carcinoma expressing progenitor cell markers. [Liver Int. 32: 430-40.](#)
6. Armstrong, A.J. *et al.* (2019) Pharmacodynamic study of radium-223 in men with bone metastatic castration resistant prostate cancer. [PLoS One. 14 \(5\): e0216934.](#)

Further Reading	1. Winter, M.J. <i>et al.</i> (2003) The epithelial cell adhesion molecule (Ep-CAM) as a morphoregulatory molecule is a tool in surgical pathology. Am J Pathol. 163 (6): 2139-48.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1870GT 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M365841:200529'

