

# Datasheet: MCA1893 BATCH NUMBER 153597

Description:	MOUSE ANTI CANINE DISTEMPER VIRUS
Specificity:	CANINE DISTEMPER VIRUS
Format:	Purified
<b>Product Type:</b>	Monoclonal Antibody
Clone:	DV2-12
Isotype:	IgG2b
Quantity:	0.25 mg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen	•			
Immunohistology - Paraffin (1)	•			
ELISA	•			
Immunoprecipitation				
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Serotec target unmasking fluid (STUF) product code

BUF025A/B/C is recommended for this purpose.

Target Species	Viral
Species Cross Reactivity	Reacts with: Raccoon dog  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.05% Sodium Azide	
Approx. Protein Concentrations	lgG concentration 1.0mg/ml	
RRID	AB_323083	

#### **Specificity**

**Mouse anti canine distemper virus antibody, clone DV2-12** recognizes canine distemper virus nucleoprotein. In western blots the antibody recognises bands of ~76 kDa, 68 kDa, 54 kDa and 32 kDa under reducing conditions.

Mouse anti canine distemper virus antibody, clone DV2-12 is useful for the detection of CDV in immunofluorescence procedures, in which the staining seen has a cytoplasmic pattern in infected cells. It is also effective in paraffin embedded material.

Mouse anti canine distemper virus antibody, clone DV2-12 is anti morbillivirus specific and reacts with phocine distemper virus, racoon distemper virus and human measles virus. It does not cross react with CAV2, CCV, CPI or CPV.

#### References

- 1. Guvenc, T. *et al.* (2008) Immunohistochemical distribution of alpha B-crystallin in the cerebellum of dogs infected with canine distemper virus. <u>Acta Vet Hung. 56: 117-23.</u>
- 2. Headley, S.A. *et al.* (2009) Diagnostic exercise: Tyzzer's disease, distemper, and coccidiosis in a pup. . Vet Pathol. 46: 151-4.
- 3. Headley, S.A. *et al.* (2009) Molecular detection of Canine distemper virus and the immunohistochemical characterization of the neurologic lesions in naturally occurring old dog encephalitis. J Vet Diagn Invest. 21: 588-97.
- 4. Bathen-Noethen, A. *et al.* (2008) Magnetic resonance imaging findings in acute canine distemper virus infection. J Small Anim Pract. 49: 460-7.
- 5. Kyöstilä, K. *et al.* (2012) A SEL1L mutation links a canine progressive early-onset cerebellar ataxia to the endoplasmic reticulum-associated protein degradation (ERAD) machinery. <u>PLoS Genet. 8 (6): e1002759.</u>
- 6. Headley, S.A. and Sukura, A. (2009) Naturally occurring systemic canine distemper virus infection in a pup Braz J Vet Pathol 2: 95-101.
- 7. Kubo, M. *et al.* (2010) Meningoencephalitis associated with Sarcocystis spp. in a free-living Japanese raccoon dog (*Nyctereutes procyonoides viverrinus*). <u>J Comp Pathol.</u> 143: 185-9.
- 8. Sánchez-Migallón Guzmán, D. *et al.* (2008) Aleutian disease serology, protein electrophoresis, and pathology of the European mink (*Mustela lutreola*) from Navarra, Spain. J Zoo Wildl Med. 39: 305-13.
- 9. Szentiks, C.A. *et al.* (2013) Polar bear encephalitis: establishment of a comprehensive next-generation pathogen analysis pipeline for captive and free-living wildlife. <u>J Comp Pathol. 2014 May;150(4):474-88.</u>

- 10. Rentería-Solís, Z. *et al.* (2014) Canine distemper outbreak in raccoons suggests pathogen interspecies transmission amongst alien and native carnivores in urban areas from Germany. Vet Microbiol. 174 (1-2): 50-9.
- 11. Woo, G.H. *et al.* (2010) Canine distemper virus infection in fennec fox (*Vulpes zerda*). J Vet Med Sci. 72 (8): 1075-9.
- 12. Spitzbarth, I. *et al.* (2010) Immunohistochemical characterization of inflammatory and glial responses in a case of necrotizing leucoencephalitis in a French bulldog. <u>J Comp</u> Pathol. 142 (2-3): 235-41.
- 13. Liang, C.T. *et al.* (2007) A non-biotin polymerized horseradish-peroxidase method for the immunohistochemical diagnosis of canine distemper. <u>J Comp Pathol. 136 (1): 57-64.</u>
- 14. Bak, W.E. and Woo, G.H. (2016) Canine Distemper in Albino Ferrets (*Mustela putorius furo*) <u>J comp Pathol</u>. 154: 121

#### **Storage**

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1893">https://www.bio-rad-antibodies.com/SDS/MCA1893</a> 10040
Regulatory	For research purposes only

## Related Products

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (STAR70...) FITC

Rabbit Anti Mouse IgG (STAR13...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

North & South America

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

## Printed on 05 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint