

Datasheet: MCA1893

BATCH NUMBER 159054

Description:	MOUSE ANTI CANINE DISTEMPER VIRUS
Specificity:	CANINE DISTEMPER VIRUS
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	DV2-12
Isotype:	IgG2b
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Serotec target unmasking fluid (STUF) product code [BUF025A/B/C](#) is recommended for this purpose.

Target Species	Viral
Species Cross Reactivity	<p>Reacts with: Raccoon dog</p> <p>N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.05% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
RRID	AB_323083
Specificity	<p>Mouse anti canine distemper virus antibody, clone DV2-12 recognizes canine distemper virus nucleoprotein. In western blots the antibody recognises bands of ~76 kDa, 68 kDa, 54 kDa and 32 kDa under reducing conditions.</p> <p>Mouse anti canine distemper virus antibody, clone DV2-12 is useful for the detection of CDV in immunofluorescence procedures, in which the staining seen has a cytoplasmic pattern in infected cells. It is also effective in paraffin embedded material.</p> <p>Mouse anti canine distemper virus antibody, clone DV2-12 is anti morbillivirus specific and reacts with phocine distemper virus, racoon distemper virus and human measles virus. It does not cross react with CAV2, CCV, CPI or CPV.</p>
References	<ol style="list-style-type: none"> Guvenc, T. <i>et al.</i> (2008) Immunohistochemical distribution of alpha B-crystallin in the cerebellum of dogs infected with canine distemper virus. Acta Vet Hung. 56: 117-23. Headley, S.A. <i>et al.</i> (2009) Diagnostic exercise: Tyzzer's disease, distemper, and coccidiosis in a pup. . Vet Pathol. 46: 151-4. Headley, S.A. <i>et al.</i> (2009) Molecular detection of Canine distemper virus and the immunohistochemical characterization of the neurologic lesions in naturally occurring old dog encephalitis. J Vet Diagn Invest. 21: 588-97. Bathen-Noethen, A. <i>et al.</i> (2008) Magnetic resonance imaging findings in acute canine distemper virus infection. J Small Anim Pract. 49: 460-7. Kyöstilä, K. <i>et al.</i> (2012) A SEL1L mutation links a canine progressive early-onset cerebellar ataxia to the endoplasmic reticulum-associated protein degradation (ERAD) machinery. PLoS Genet. 8 (6): e1002759. Headley, S.A. and Sukura, A. (2009) Naturally occurring systemic canine distemper virus infection in a pup Braz J Vet Pathol 2: 95-101. Kubo, M. <i>et al.</i> (2010) Meningoencephalitis associated with Sarcocystis spp. in a free-living Japanese raccoon dog (<i>Nyctereutes procyonoides viverrinus</i>). J Comp Pathol. 143: 185-9. Sánchez-Migallón Guzmán, D. <i>et al.</i> (2008) Aleutian disease serology, protein electrophoresis, and pathology of the European mink (<i>Mustela lutreola</i>) from Navarra, Spain. J Zoo Wildl Med. 39: 305-13. Szentiks, C.A. <i>et al.</i> (2013) Polar bear encephalitis: establishment of a comprehensive next-generation pathogen analysis pipeline for captive and free-living wildlife. J Comp Pathol. 2014 May;150(4):474-88.

10. Rentería-Solís, Z. *et al.* (2014) Canine distemper outbreak in raccoons suggests pathogen interspecies transmission amongst alien and native carnivores in urban areas from Germany. [Vet Microbiol. 174 \(1-2\): 50-9.](#)
11. Woo, G.H. *et al.* (2010) Canine distemper virus infection in fennec fox (*Vulpes zerda*). [J Vet Med Sci. 72 \(8\): 1075-9.](#)
12. Spitzbarth, I. *et al.* (2010) Immunohistochemical characterization of inflammatory and glial responses in a case of necrotizing leucoencephalitis in a French bulldog. [J Comp Pathol. 142 \(2-3\): 235-41.](#)
13. Liang, C.T. *et al.* (2007) A non-biotin polymerized horseradish-peroxidase method for the immunohistochemical diagnosis of canine distemper. [J Comp Pathol. 136 \(1\): 57-64.](#)
14. Bak, W.E. and Woo, G.H. (2016) Canine Distemper in Albino Ferrets (*Mustela putorius furo*) [J comp Pathol. 154: 121](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1893>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
 Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
 Goat Anti Mouse IgG (STAR76...) [RPE](#)
 Goat Anti Mouse IgG (STAR70...) [FITC](#)
 Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
 Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
 Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
 Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
 Goat Anti Mouse IgG (STAR77...) [HRP](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

