

## Datasheet: MCA1895

**BATCH NUMBER 166352**

<b>Description:</b>	MOUSE ANTI DOG IgE
<b>Specificity:</b>	IgE
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	E6-2A1
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.25 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Dog
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	<0.1% sodium azide (NaN <sub>3</sub> )
<b>Approx. Protein</b>	IgG concentration 1.0 mg/ml

## Concentrations

**Immunogen** Affinity purified IgE preparation from heavily parasitized and allergic dog serum.

**RRID** AB\_323084

**Specificity** **Mouse anti Dog IgE antibody, clone E6-2A1** recognises canine IgE. E6-2A1 does not cross react with dog IgM, IgA, IgG1 or IgG2. Western blot analysis against affinity purified dog IgE using Mouse anti Dog IgE clone E6-2A1 demonstrates a single major band of ~62-65 kDa under reducing conditions.

**ELISA** Mouse anti Dog IgE antibody, clone E6-2A1 may be used as a capture antibody in a sandwich ELISA in combination with [MCA1896B](#) as the detection antibody.

**References**

1. Lin, T.Y. *et al.* (2006a) Generation and characterization of bone marrow-derived cultured canine mast cells. [Vet Immunol Immunopathol. 113 \(1-2\): 37-52.](#)
2. Lin, T.Y. & London, C.A. (2006b) A functional comparison of canine and murine bone marrow derived cultured mast cells. [Vet Immunol Immunopathol. 114 \(3-4\): 320-34.](#)
3. Khantavee, N. *et al.* (2020) Antibody levels to *Malassezia pachydermatis*. and *Staphylococcus pseudintermedius*. in atopic dogs and their relationship with lesion scores. [Vet Dermatol. 31 \(2\): 111-115.](#)
4. Khantavee, N. *et al.* (2021) Immunoglobulin G1 subclass responses can be used to detect specific allergy to the house dust mites *Dermatophagoides farinae*. and *Dermatophagoides pteronyssinus*. in atopic dogs. [BMC Vet Res. 17 \(1\): 71.](#)
5. Mason, K. *et al.* (2023) Immunoglobulin E-specific allergens against leaf in serum of dogs with clinical features of grass leaf allergy. [Vet Dermatol. 34 \(5\): 393-403.](#)

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

**Guarantee** 12 months from date of despatch

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1895>  
10040

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)

Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),  
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),  
[FITC](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

### **Recommended Negative Controls**

[MOUSE IgG2a NEGATIVE CONTROL \(MCA929\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M415520:230105'

**Printed on 11 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)