

## Datasheet: MCA1898PE

<b>Description:</b>	RAT ANTI MOUSE CD115:RPE
<b>Specificity:</b>	CD115
<b>Other names:</b>	c-fms
<b>Format:</b>	RPE
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	604B5 2E11
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse		
<b>Product Form</b>	Purified IgG - lyophilized		
<b>Reconstitution</b>	Reconstitute with 1.0 ml distilled water		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	RPE 488nm laser	496	578
<b>Preparation</b>	Purified IgG prepared by ion exchange chromatography		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative Stabilisers</b>	0.09% Sodium Azide 1% Bovine Serum Albumin		
<b>Immunogen</b>	Raw 264 cells.		

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P09581</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">12978</a> Csf1r <a href="#">Related reagents</a>
<b>Synonyms</b>	Csfmr, Fms
<b>RRID</b>	AB_323314
<b>Fusion Partners</b>	Spleen cells from immunised Sprague Dawley rats were fused with cells of the P3-653 myeloma cell line.
<b>Specificity</b>	<p><b>Rat anti Mouse CD115 antibody, clone 604B5 2E11</b> recognizes the murine CD115 cell surface antigen, also known as the M-CSF receptor and as c-fms. Murine CD115 is a 958 amino acid 110kDa single pass type I transmembrane glycoprotein with tyrosine kinase activity and acts as a membrane receptor for M-CSF.</p> <p>CD115 is expressed by cells of the monocytic lineage and by progenitor cells. Rat anti Mouse CD115 antibody, clone 604B5 2E11 has been shown to inhibit <i>in vitro</i> colony formation in response to M-CSF in both rats and mice.</p>
<b>Flow Cytometry</b>	<p>Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<a href="#">BUF041A/B</a>).</p>
<b>References</b>	<ol style="list-style-type: none"> <li>Shadduck, R.K. <i>et al.</i> (1996) Paradoxical stimulation of normal and leukemic rat hematopoiesis by monoclonal antibody to CSF-1 receptor. <a href="#">Exp Hematol. 24 (2): 314-7.</a></li> <li>Gilmore, G.L. &amp; Shadduck, R.K. (1995) Inhibition of day-12 spleen colony-forming units by a monoclonal antibody to the murine macrophage/monocyte colony-stimulating factor receptor. <a href="#">Blood. 85 (10): 2731-4.</a></li> <li>MacDonald, K.P. <i>et al.</i> (2005) The colony-stimulating factor 1 receptor is expressed on dendritic cells during differentiation and regulates their expansion. <a href="#">J Immunol. 175 (3): 1399-405.</a></li> <li>Buono, M. <i>et al.</i> (2010) Sulfatase modifying factor 1-mediated fibroblast growth factor signaling primes hematopoietic multilineage development. <a href="#">J Exp Med. 207 (8): 1647-60.</a></li> <li>Weingärtner, O. <i>et al.</i> (2011) Differential effects on inhibition of cholesterol absorption by plant stanol and plant sterol esters in apoE<sup>-/-</sup> mice. <a href="#">Cardiovasc Res. 90: 484-92.</a></li> <li>Hayashi, Y. <i>et al.</i> (2010) Lumican is required for neutrophil extravasation following corneal injury and wound healing. <a href="#">J Cell Sci. 123: 2987-95.</a></li> <li>Herold, S. <i>et al.</i> (2008) Lung epithelial apoptosis in influenza virus pneumonia: the role of macrophage-expressed TNF-related apoptosis-inducing ligand. <a href="#">J Exp Med. 205: 3065-77.</a></li> <li>Pillai, M.M. <i>et al.</i> (2009) Inducible transgenes under the control of the hCD68 promoter identifies mouse macrophages with a distribution that differs from the F4/80 - and CSF-1R-expressing populations. <a href="#">Exp Hematol. 37: 1387-92.</a></li> </ol>

9. Sasmono, R.T. *et al.* (2007) Mouse neutrophilic granulocytes express mRNA encoding the macrophage colony-stimulating factor receptor (CSF-1R) as well as many other macrophage-specific transcripts and can transdifferentiate into macrophages *in vitro* in response to CSF-1. [J Leukoc Biol. 82: 111-23.](#)
10. Dioszeghy V *et al.* (2008) 12/15-Lipoxygenase regulates the inflammatory response to bacterial products *in vivo*. [J Immunol. 181 \(9\): 6514-24.](#)
11. Fancke B *et al.* (2008) M-CSF: a novel plasmacytoid and conventional dendritic cell poietin. [Blood. 111 \(1\): 150-9.](#)
12. van Dommelen, S.L. *et al.* (2006) Perforin and granzymes have distinct roles in defensive immunity and immunopathology. [Immunity. 25 \(5\): 835-48.](#)
13. Scheller, M. *et al.* (1999) Altered development and cytokine responses of myeloid progenitors in the absence of transcription factor, interferon consensus sequence binding protein. [Blood. 94 \(11\): 3764-71.](#)
14. O'Dea, K.P. *et al.* (2009) Mobilization and margination of bone marrow Gr-1high monocytes during subclinical endotoxemia predisposes the lungs toward acute injury. [J Immunol. 182 \(2\): 1155-66.](#)
15. Nahrendorf, M. *et al.* (2007) The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. [J Exp Med. 204 \(12\): 3037-47.](#)
16. Juss, J.K. *et al.* (2012) Functional redundancy of class I phosphoinositide 3-kinase (PI3K) isoforms in signaling growth factor-mediated human neutrophil survival. [PLoS One. 7 \(9\): e45933.](#)

---

**Storage** Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #20487 available at: 20487: <https://www.bio-rad-antibodies.com/uploads/MSDS/20487.pdf>

---

**Regulatory** For research purposes only

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M378570:210222'

**Printed on 08 Oct 2021**