

Datasheet: MCA1899F

Description:	MOUSE ANTI HORSE PAN B-CELLS:FITC
Specificity:	PAN B-CELLS
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CVS36
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
ELISA	▪			
Immunofluorescence			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Horse						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant						
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin						
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml						

Immunogen	Purified Equine Ig.
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Horse Pan B-Cells, clone CVS36 is a monoclonal antibody directed against equine Ig light chains. Characterization studies demonstrated that clone CVS36 bound 100% of CD5-ve peripheral blood lymphocytes and recognized all equine B-cells (Lunn <i>et al.</i> 1998) As the antigen recognized by clone CVS36 appears to be present on the surface of all equine B-cells it is therefore is a reagent that can be used as a pan B-cell marker for domestic horses (Breathnach <i>et al.</i> 2005).</p> <p>Specific anti equine reagents have yet to be fully characterized for the typically recognized B-cell makers such as CD19, CD20, CD21, CD22 and CD79. While testing has demonstrated the cross reactivity of some monoclonal B-cell markers raised against other species with equine B cells, such as the Mouse anti Human CD79a antibody, (clone HM57), an overview of which may be found in the report of the second equine leucocyte antigen workshop (Lunn <i>et al.</i> 1998).</p> <p>With specificity for equine Ig light chains, clone CVS36 may be used to detect all equine immunoglobulin classes and subclasses in ELISA applications (Lunn <i>et al.</i> 1998).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul
References	<ol style="list-style-type: none"> Schneider, R. (2010) Analysis of antibody subtypes and T lymphocyte activation following vaccination of healthy foals against West Nile Virus In M.Sc thesis University of Pennsylvania, Chapter 3, p.34-45 Lunn, D.P. <i>et al.</i> (1998) Report of the Second Equine Leucocyte Antigen Workshop, Squaw valley, California, July 1995. Vet Immunol Immunopathol. 62 (2): 101-43. Umlauf, C. (2004) Herstellung und Charakterisierung monoklonaler Antikörper gegen equine Leukozyten In Phd thesis Ludwig-Maximilians-Universität München Tomlinson, J.E. <i>et al.</i> (2018) Multispectral fluorescence-activated cell sorting of B and T cell subpopulations from equine peripheral blood. Vet Immunol Immunopathol. 199: 22-31.
Further Reading	<ol style="list-style-type: none"> Lunn, D.P. <i>et al.</i> (1991) Three monoclonal antibodies identifying antigens on all equine T lymphocytes, and two mutually exclusive T-lymphocyte subsets. Immunology. 74 (2): 251-7. Sheoran, A.S. <i>et al.</i> (1998) Monoclonal antibodies to subclass-specific antigenic determinants on equine immunoglobulin gamma chains and their characterization. Vet Immunol Immunopathol. 62 (2): 153-65.
Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[MOUSE ANTI HORSE PAN B-CELLS \(MCA1899GA\)](#)

[MOUSE ANTI HORSE PAN B-CELLS:RPE \(MCA1899PE\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M383182:210513'

Printed on 06 Jan 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)