

Datasheet: MCA189B BATCH NUMBER 160192

Description:	MOUSE ANTI RAT IgM HEAVY CHAIN:Biotin
Specificity:	IgM HEAVY CHAIN
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	MARM-4
Isotype:	lgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	•			125ng/ml as detecting antibody

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Rat
Purified IgG conjugated to Biotin - liquid
Purified IgG prepared by affinity chromatography from tissue culture supernatant
Phosphate buffered saline
<0.1% Sodium Azide (NaN ₃) 50% Glycerol
IgG concentration 1.0 mg/ml

Immunogen	IR202 rat IgM myeloma protein
RRID	AB_321875
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	Mouse anti Rat IgM Heavy Chain antibody, clone MARM-4 recognizes the mu heavy chain of rat immunoglobulin and does not cross-react with other immunoglobulin classes.
References	1. Bazin, H. <i>et al.</i> (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. Eur J Immunol. 4 (1): 44-8. 2. Bazin, H. <i>et al.</i> (1978) Transplantable IgD immunoglobulin-secreting tumors in rat. J Immunol. 121 (5): 2077-82. 3. Khalifeh, M.S. <i>et al.</i> (2010) Investigation ofthe role of tumour necrosis factor-{alpha}, interleukin-1{beta}, interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. Lab Anim. 44: 143-9. 4. Bézie, S. <i>et al.</i> (2015) Compensatory Regulatory Networks between CD8 T, B, and Myeloid Cells in Organ Transplantation Tolerance. J Immunol. 195 (12): 5805-15. 5. Ueta, H. <i>et al.</i> (2018) Single blood transfusion induces the production of donor-specific alloantibodies and regulatory T cells mainly in the spleen. Int Immunol. 30 (2): 53-67. 6. Tulinská, J. <i>et al.</i> (2018) Humoral and cellular immune response in Wistar Han RCC rats fed two genetically modified maize MON810 varieties for 90 days (EU 7th Framework Programme project GRACE). Arch Toxicol. 92 (7): 2385-99.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/MCA189B 10049
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:Biotin (MCA1209B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Worldwide

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint