

### Datasheet: MCA189F

Description:	MOUSE ANTI RAT IgM HEAVY CHAIN:FITC
Specificity:	IgM HEAVY CHAIN
Format:	FITC
<b>Product Type:</b>	Monoclonal Antibody
Clone:	MARM-4
Isotype:	lgG1
Quantity:	0.5 mg

#### **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			10ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat			
Product Form	Purified IgG conjugate	1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nn	n)
	FITC	490	525	
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant			
Buffer Solution	Phosphate buffered saline			
Preservative	0.1% Sodium Azide			
Stabilisers	50% Glycerol			
Approx. Protein Concentrations	IgG concentration 1.0	mg/ml		
Immunogen	IR202 rat IgM myelom	na protein		

RRID	AB_321876
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	Mouse anti Rat IgM Heavy Chain antibody, clone MARM-4 recognizes the mu heavy chain of rat immunoglobulin and does not cross-react with other immunoglobulin classes.
Flow Cytometry	Use 50ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	1. Bazin, H. <i>et al.</i> (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. <u>Eur J Immunol. 4 (1): 44-8.</u> 2. Bazin, H. <i>et al.</i> (1978) Transplantable IgD immunoglobulin-secreting tumors in rat. <u>J Immunol. 121 (5): 2077-82.</u> 3. Khalifeh, M.S. <i>et al.</i> (2010) Investigation ofthe role of tumour necrosis factor-{alpha}, interleukin-1{beta}, interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. <u>Lab Anim. 44: 143-9.</u> 4. Bézie, S. <i>et al.</i> (2015) Compensatory Regulatory Networks between CD8 T, B, and Myeloid Cells in Organ Transplantation Tolerance. <u>J Immunol. 195 (12): 5805-15.</u> 5. Ueta, H. <i>et al.</i> (2018) Single blood transfusion induces the production of donor-specific alloantibodies and regulatory T cells mainly in the spleen. <u>Int Immunol. 30 (2): 53-67.</u> 6. Tulinská, J. <i>et al.</i> (2018) Humoral and cellular immune response in Wistar Han RCC rats fed two genetically modified maize MON810 varieties for 90 days (EU 7th Framework Programme project GRACE). <u>Arch Toxicol. 92 (7): 2385-99.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10328 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA189F">https://www.bio-rad-antibodies.com/SDS/MCA189F</a> 10328
Regulatory	For research purposes only

# **Related Products**

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA1209F)

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_us@bio-rad.com

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint