

Datasheet: MCA189F

Description:	MOUSE ANTI RAT IgM HEAVY CHAIN:FITC
Specificity:	IgM HEAVY CHAIN
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MARM-4
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			10ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.1% Sodium Azide		
Stabilisers	50% Glycerol		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	IR202 rat IgM myeloma protein		
RRID	AB_321876		
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.		

Specificity	Mouse anti Rat IgM Heavy Chain antibody, clone MARM-4 recognizes the mu heavy chain of rat immunoglobulin and does not cross-react with other immunoglobulin classes.
Flow Cytometry	Use 50ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Bazin, H. <i>et al.</i> (1974) Three classes and four (sub)classes of rat immunoglobulins: IgM, IgA, IgE and IgG1, IgG2a, IgG2b, IgG2c. Eur J Immunol. 4 (1): 44-8. 2. Bazin, H. <i>et al.</i> (1978) Transplantable IgD immunoglobulin-secreting tumors in rat. J Immunol. 121 (5): 2077-82. 3. Khalifeh, M.S. <i>et al.</i> (2010) Investigation of the role of tumour necrosis factor-α, interleukin-1β, interleukin-10, nitric oxide and rheumatoid factor-immunoglobulin M in a rat model of arthritis. Lab Anim. 44: 143-9. 4. Bézie, S. <i>et al.</i> (2015) Compensatory Regulatory Networks between CD8 T, B, and Myeloid Cells in Organ Transplantation Tolerance. J Immunol. 195 (12): 5805-15. 5. Ueta, H. <i>et al.</i> (2018) Single blood transfusion induces the production of donor-specific alloantibodies and regulatory T cells mainly in the spleen. Int Immunol. 30 (2): 53-67.
Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10328 available at: 10328: https://www.bio-rad-antibodies.com/uploads/MSDS/10328.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA1209F\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M365878:200529'

Printed on 11 Aug 2020