

## Datasheet: MCA1901GA

**BATCH NUMBER 154526**

<b>Description:</b>	MOUSE ANTI HORSE IgGb
<b>Specificity:</b>	IgGb
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	CVS39
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/5000 - 1/50000
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Horse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified equine IgGb.
<b>Fusion Partners</b>	Spleen cells from immunised Balb/c mice were fused with cells of the mouse X63-Ag8.653 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Horse IgGb antibody, clone CVS39</b> recognizes equine IgGb, recently classified as equine IgG4/7 (<a href="#">Keggan et al. 2013</a>), and shows no significant binding to other IgG subclasses. Equine IgGb is an immunoglobulin with a molecular weight (MW) of approximately 160kDa, the MW of the light chain being 27kDa, common to all equine IgG subclasses, while the heavy chain has a MW of 53kDa, specific to IgGb (<a href="#">Sheoran et al. 1998</a>). Equine IgGb has been demonstrated to be involved in long lasting immunity following respiratory-related viral infections such as Equine Herpes virus Type 4 (EHV-4) (<a href="#">Mizukoshi et al. 2002</a>) and Equine Influenza (<a href="#">Soboll et al. 2003</a>).</p> <p>In addition to Mouse anti Horse IgGb, clone CVS39, Bio-Rad also offer a range of other monoclonal and polyclonal antibodies that are specific to equine immunoglobulin classes and subclasses.</p>
<b>References</b>	<ol style="list-style-type: none"> <li>1. Sheoran, A.S. <i>et al.</i> (1998) Monoclonal antibodies to subclass-specific antigenic determinants on equine immunoglobulin gamma chains and their characterization. <a href="#">Vet Immunol Immunopathol. 62 (2): 153-65.</a></li> <li>2. Lewis, M.J. <i>et al.</i> (2008) The different effector function capabilities of the seven equine IgG subclasses have implications for vaccine strategies. <a href="#">Mol Immunol. 45 (3): 818-27.</a></li> <li>3. Keggan, A. <i>et al</i> (2013) Production of seven monoclonal equine immunoglobulins isotyped by multiplex analysis <a href="#">Vet Immunol Immunopathol.153(3-4):187-93</a></li> </ol>
<b>Further Reading</b>	<ol style="list-style-type: none"> <li>1. Mizukoski, F. <i>et al</i> (2002) IgG antibody subclass response against equine herpesvirus type 4 in horses. <a href="#">Vet Immunol Immunopathol. 88(1-2):97-101.</a></li> <li>2. Soboll, G. <i>et al</i> (2003) Regional antibody and cellular immune responses to equine influenza virus infection, and particle mediated DNA vaccination. <a href="#">Vet Immunol Immunopathol. 94(1-2):47-62.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1901GA">https://www.bio-rad-antibodies.com/SDS/MCA1901GA</a> 10040

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Useful Reagents

[PURIFIED HORSE IgG \(PEP001\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M365882:200529'

Printed on 18 Jan 2024