

Datasheet: MCA191

BATCH NUMBER 161118

Description:	MOUSE ANTI RAT IgA HEAVY CHAIN
Specificity:	IgA HEAVY CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MARA-1
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% sodium azide

Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Purified IR1060 IgA rat myeloma protein.
RRID	AB_322196
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	Mouse anti Rat IgA Heavy Chain antibody, clone MARA-1 recognizes the alpha heavy chain of rat immunoglobulin. Mouse anti Rat IgA Heavy Chain antibody, clone MARA-1 shows no cross-reactivity with other rat immunoglobulin classes.
ELISA	Mouse anti Rat IgA Heavy Chain antibody, clone MARA-1 may be used as a coating antibody in a sandwich ELISA in combination with Mouse anti Rat κ/λ antibody, clone MARK-1/MARL-15 (MCA1296P) as detection reagent and purified rat IgA (PRP01) as antigen..
References	<ol style="list-style-type: none"> 1. Bjersing, J.L. <i>et al.</i> (2002) Loss of ileal IgA+ plasma cells and of CD4+ lymphocytes in ileal Peyer's patches of vitamin A deficient rats. Clin Exp Immunol. 130: 404-8. 2. Budeč, M. <i>et al.</i> (2007) Possible mechanism of acute effect of ethanol on intestinal IgA expression in rat. Int Immunopharmacol. 7: 858-63. 3. Budeč, M. <i>et al.</i> (2009) Blockade of nitric oxide synthesis modulates rat immunoglobulin A. Neuroimmunomodulation. 16: 155-61. 4. Hahn, A. <i>et al.</i> (2010) Mesenteric lymph nodes are not required for an intestinal immunoglobulin A response to oral cholera toxin. Immunology. 129: 427-36. 5. Herías, M.V. <i>et al.</i> (1999) Immunomodulatory effects of Lactobacillus plantarum colonizing the intestine of gnotobiotic rats Clin Exp Immunol. 116: 283-90. 6. Ito, H. <i>et al.</i> (2011) Degree of polymerization of inulin-type fructans differentially affects number of lactic acid bacteria, intestinal immune functions, and immunoglobulin A secretion in the rat cecum. J Agric Food Chem. 59 (10): 5771-8. 7. Kushnir, N. <i>et al.</i> (1998) Dendritic cells and resting B cells form clusters in vitro and in vivo: T cell independence, partial LFA-1 dependence, and regulation by cross-linking surface molecules. J Immunol. 160: 1774-81. 8. Goodrich, M.E. and McGee, D.W. (1998) Regulation of mucosal B cell immunoglobulin secretion by intestinal epithelial cell-derived cytokines. Cytokine. 10: 948-55. 9. Heel, K.A. <i>et al.</i> (1998) The effect of minimum luminal nutrition on mucosal cellularity and immunity of the gut. J Gastroenterol Hepatol. 13: 1015-9. 10. Pérez-Cano FJ (2005) Neonatal immunoglobulin secretion and lymphocyte phenotype in rat small intestine lamina propria. Pediatr Res. 58: 164-9. 11. Peng, Z. <i>et al.</i> (1996) Cross-reactivity and molecular mass of the epsilon chains of the IgE antibodies in dogs, humans, rats, and mice. Int Arch Allergy Immunol. 110: 149-55. 12. Nayak, B.N. <i>et al.</i> (2009) Energy-restricted diets result in higher numbers of CD4+, CD8+, immunoglobulins (A, M, and G), and CD45RA cells in spleen and CD4+, immunoglobulin A, and CD45RA cells in colonic lamina propria of rats. Nutr Res. 2009 Jul;29(7):487-93. 13. Komura, M. <i>et al.</i> (2014) A short-term ingestion of fructo-oligosaccharides increases

immunoglobulin A and mucin concentrations in the rat cecum, but the effects are attenuated with the prolonged ingestion. [Biosci Biotechnol Biochem. 78: 1592-602.](#)

14. Ito, H. *et al.* (2011) Degree of polymerization of inulin-type fructans differentially affects number of lactic acid bacteria, intestinal immune functions, and immunoglobulin A secretion in the rat cecum. [J Agric Food Chem. 59: 5771-8.](#)

15. Hino, S. *et al.* (2020) Mucin-Derived O-Glycans Act as Endogenous Fiber and Sustain Mucosal Immune Homeostasis via Short-Chain Fatty Acid Production in Rat Cecum. [J Nutr. 150 \(10\): 2656-65.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA19110040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M365887:200529'

Printed on 12 Aug 2023

