

Datasheet: MCA193

BATCH NUMBER 148596

Description:	MOUSE ANTI RAT IgE
Specificity:	IgE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MARE-1
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by immunoaffinity on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide
Approx. Protein	IgG concentration 1.0 mg/ml

Concentrations

Immunogen Rat IR162, IR1016, IR2 and IR410 IgE myeloma proteins.

External Database Links

UniProt:

[P01855](#) [Related reagents](#)

Entrez Gene:

[299351](#) Ighe [Related reagents](#)

RRID AB_321899

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.

Specificity **Mouse anti Rat IgE antibody, clone MARE-1** recognizes rat epsilon heavy chain of immunoglobulin IgE and does not cross-react with other classes of rat immunoglobulin.

Mouse anti Rat IgE antibody, clone MARE-1 binds to rat IgE with an avidity of $4 \times 10^9 M^{-1}$

ELISA This antibody may be used as a coating antibody in a sandwich ELISA in combination with clone MARK-1/MARL-15 (Product code MCA1296P) as detection reagent and purified rat IgE (Product code PRP07A).

References

1. Negrão-Corrêa, D. *et al.* (1996) Intestinal transport and catabolism of IgE: a major blood-independent pathway of IgE dissemination during a *Trichinella spiralis* infection of rats. [J Immunol. 157 \(9\): 4037-44.](#)
2. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. I. Rapid purification from *in vitro* culture supernatants. [J Immunol Methods. 66 \(2\): 261-9.](#)
3. Bazin, H. *et al.* (1974) Transplantable immunoglobulin-secreting tumours in rats. IV. Sixty-three IgE-secreting immunocytoma tumours. [Immunology. 26 \(4\): 713-23.](#)
4. Bazin, H. *et al.* (1978) Transplantable IgD immunoglobulin-secreting tumors in rat. [J Immunol. 121 \(5\): 2077-82.](#)
5. Bazin, H. *et al.* (1984) Rat monoclonal antibodies. II. A rapid and efficient method of purification from ascitic fluid or serum. [J Immunol Methods. 71 \(1\): 9-16.](#)
6. Cho, J.K. & Cho, S.W. (2000) Shared epitope for monoclonal IR162 between *Anisakis simplex* larvae and *Clonorchis sinensis* and cross-reactivity between antigens. [J Parasitol. 86 \(5\): 1145-9.](#)
7. Silveira, M.R. *et al.* (2002) Infection with *Strongyloides venezuelensis* induces transient airway eosinophilic inflammation, an increase in immunoglobulin E, and hyperresponsiveness in rats. [Infect Immun. 70: 6263-72.](#)
8. Korinek, M. *et al.* (2016) Anti-allergic potential of *Typhonium blumei*: inhibition of degranulation via suppression of PI3K/PLC γ 2 phosphorylation and calcium influx. [Phytomedicine. 23 \(14\): 1706-15.](#)
9. Bąbolewska, E. & Brzezińska-błaszczyk, E. (2015) Human-derived cathelicidin LL-37 directly activates mast cells to proinflammatory mediator synthesis and migratory response. [Cell Immunol. 293 \(2\): 67-73.](#)

10. Agier, J. *et al.* (2018) Cathelicidin LL-37 Affects Surface and Intracellular Toll-Like Receptor Expression in Tissue Mast Cells. [J Immunol Res. 2018: 7357162.](#)
11. Ueta, H. *et al.* (2018) Single blood transfusion induces the production of donor-specific alloantibodies and regulatory T cells mainly in the spleen. [Int Immunol. 30 \(2\): 53-67.](#)
12. Witczak, P. *et al.* (2020) The Response of Tissue Mast Cells to TLR3 Ligand Poly(I:C) Treatment. [J Immunol Res. 2020: 2140694.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA19310040>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M365912:200529'

Printed on 22 Oct 2024