

Datasheet: MCA1940A488

BATCH NUMBER 1112

Description:	MOUSE ANTI HUMAN CD19:Alexa Fluor® 488			
Specificity:	CD19			
Format:	ALEXA FLUOR® 488			
Product Type:	Monoclonal Antibody			
Clone:	LT19			
Isotype:	lgG1			
Quantity:	100 TESTS/1ml			

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human			
Product Form	Purified IgG conjuga			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nr	m)
	Alexa Fluor®488	495	519	
Preparation Buffer Solution	Phosphate buffered	change chromatography.		
Preservative	0.09% Sodium Azid	е		
Stabilisers	1% Bovine Serur			
Approx. Protein Concentrations	IgG concentration 0	.05 mg/ml		

External Database Links

UniProt:

P15391 Related reagents

Entrez Gene:

930 CD19 Related reagents

RRID

AB 324914

Specificity

Mouse anti Human CD19 antibody, clone LT19 recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells (de Rie *et al.* 1989).

CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 (<u>Bradbury et al. 1992</u>), CD9, CD81 and CD82 (<u>Horváth et al. 1998</u>). CD19 is implicated in the down-regulation of B cell growth and proliferation (<u>Pezzutto et al. 1987</u>).

Flow Cytometry

Use 10ul of the suggested working dilution to label 1 x 10⁶ cells in 100ul

References

- 1. Hughes, G.J. *et al.* (2007) Virus immunocapture provides evidence of CD8 lymphocytederived HIV-1 *in vivo*. AIDS. 21: 1507-13.
- 2. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. <u>Diabetes</u>. 58: 138-45.
- 3. McIntosh, K. *et al.* (2006) The immunogenicity of human adipose-derived cells: temporal changes *in vitro*. <u>Stem Cells. 24: 1246-53.</u>
- 4. Sengstake, S. *et al.* (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. Int Immunol. 18 (7): 1171-8.
- 5. Villarroel Dorrego, M. *et al.* (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. <u>Br J Dermatol. 154: 231-8.</u>
- 6. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. <u>Blood. 118: 348-57.</u>
- 7. Lacal, P.M. *et al.* (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. <u>J Pharmacol Exp Ther. 347:</u> 164-72.
- 8. Franz, B. *et al.* (2011) Ex vivo characterization and isolation of rare memory B cells with antigen tetramers. <u>Blood. 118: 348-57.</u>
- 9. Girbl, T. *et al.* (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. <u>Cancer Res. 73: 561-70.</u>
- 10. Hertzberg, L. *et al.* (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. <u>Blood. 115: 1006-17.</u>
- 11. Kakko, T. et al. (2011) Inflammatory effects of blood leukocytes: association with

vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. <u>Diab</u> Vasc Dis Res. 8: 221-8.

- 12. Dorvignit, D. *et al.* (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. MAbs. 4 (4): 488-96.
- 13. Karlsen, M. *et al.* (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. <u>Scand J Immunol. 82 (6): 523-31.</u>
- 14. Clark, L.E. *et al.* (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. <u>Nat Commun. 9 (1): 1884.</u>
- 15. Gu, Y. *et al.* (2019) Defining the structural basis for human alloantibody binding to human leukocyte antigen allele HLA-A*11:01. <u>Nat Commun. 10 (1): 893.</u>
- 16. Yang, C. *et al.* (2013) B cells promote tumor progression via STAT3 regulated-angiogenesis. PLoS One. 8 (5): e64159.

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Acknowledgements

This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1940A488 10041

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL: Alexa Fluor® 488 (MCA928A488)

Recommended Useful Reagents

<u>HUMAN SEROBLOCK (BUF070A)</u> <u>HUMAN SEROBLOCK (BUF070B)</u>

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365928:200529'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint