

Datasheet: MCA1940A488T

Description:	MOUSE ANTI HUMAN CD19:Alexa Fluor® 488
Specificity:	CD19
Format:	ALEXA FLUOR® 488
Product Type:	Monoclonal Antibody
Clone:	LT19
Isotype:	IgG1
Quantity:	25 TESTS/0.25ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Alexa Fluor® 488 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®488	495	519
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN ₃)		
Stabilisers	1% bovine serum albumin		
Approx. Protein Concentrations	IgG concentration 0.05 mg/ml		

External Database
Links

UniProt:

[P15391](#) [Related reagents](#)

Entrez Gene:

[930](#) CD19 [Related reagents](#)

RRID

AB_1101040

Specificity

Mouse anti Human CD19 antibody, clone LT19 recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells ([de Rie *et al.* 1989](#)).

CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 ([Bradbury *et al.* 1992](#)), CD9, CD81 and CD82 ([Horváth *et al.* 1998](#)). CD19 is implicated in the down-regulation of B cell growth and proliferation ([Pezzutto *et al.* 1987](#)).

Flow Cytometry

Use 10µl of the suggested working dilution to label 1 x 10⁶ cells in 100µl

References

1. Hughes, G.J. *et al.* (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 *in vivo*. [AIDS. 21: 1507-13.](#)
2. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. [Diabetes. 58: 138-45.](#)
3. McIntosh, K. *et al.* (2006) The immunogenicity of human adipose-derived cells: temporal changes *in vitro*. [Stem Cells. 24: 1246-53.](#)
4. Sengstake, S. *et al.* (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. [Int Immunol. 18 \(7\): 1171-8.](#)
5. Villarreal Dorrego, M. *et al.* (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. [Br J Dermatol. 154: 231-8.](#)
6. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
7. Lacal, P.M. *et al.* (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. [J Pharmacol Exp Ther. 347: 164-72.](#)
8. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
9. Girbl, T. *et al.* (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. [Cancer Res. 73: 561-70.](#)
10. Hertzberg, L. *et al.* (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. [Blood. 115: 1006-17.](#)
11. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with

vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab Vasc Dis Res. 8: 221-8.](#)

12. Dorvignit, D. *et al.* (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. [MAbs. 4 \(4\): 488-96.](#)

13. Karlsen, M. *et al.* (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. [Scand J Immunol. 82 \(6\): 523-31.](#)

14. Clark, L.E. *et al.* (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. [Nat Commun. 9 \(1\): 1884.](#)

15. Gu, Y. *et al.* (2019) Defining the structural basis for human alloantibody binding to human leukocyte antigen allele HLA-A*11:01. [Nat Commun. 10 \(1\): 893.](#)

16. Yang, C. *et al.* (2013) B cells promote tumor progression via STAT3 regulated-angiogenesis. [PLoS One. 8 \(5\): e64159.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Acknowledgements This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1940A488T>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 488 \(MCA928A488\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M421898:230808'

Printed on 19 Oct 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)