

## Datasheet: MCA1940A700T

|                      |  |
|----------------------|--|
| <b>Description:</b>  | MOUSE ANTI HUMAN CD19:Alexa Fluor® 700 |
| <b>Specificity:</b>  | CD19                                   |
| <b>Format:</b>       | ALEXA FLUOR® 700                       |
| <b>Product Type:</b> | Monoclonal Antibody                    |
| <b>Clone:</b>        | LT19                                   |
| <b>Isotype:</b>      | IgG1                                   |
| <b>Quantity:</b>     | 25 TESTS/0.25ml                        |

## Product Details

**RRID** AB\_1101045

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ■   |    |                | Neat - 1/10        |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Human

**Product Form** Purified IgG conjugated to Alexa Fluor® 700 - liquid

| Max Ex/Em | Fluorophore     | Excitation Max (nm) | Emission Max (nm) |
|-----------|-----------------|---------------------|-------------------|
|           | Alexa Fluor®700 | 702                 | 723               |

**Preparation** Purified IgG from tissue culture supernatant prepared by Protein A chromatography.

**Buffer Solution** Phosphate buffered saline

**Preservative** 0.09% Sodium Azide  
**Stabilisers** 1% Bovine Serum Albumin

**Approx. Protein Concentrations** IgG concentration 0.05 mg/ml

**External Database Links**  
**UniProt:**  
[P15391](https://www.uniprot.org/entry/P15391) [Related reagents](#)

**Entrez Gene:**

---

|                       |  |
|-----------------------|--|
| <b>Specificity</b>    | <p><b>Mouse anti Human CD19 antibody, clone LT19</b> recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells (<a href="#">de Rie et al. 1989</a>).</p> <p>CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 (<a href="#">Bradbury et al. 1992</a>), CD9, CD81 and CD82 (<a href="#">Horváth et al. 1998</a>). CD19 is implicated in the down-regulation of B cell growth and proliferation (<a href="#">Pezzutto et al. 1987</a>).</p>  |
| <b>Flow Cytometry</b> | Use 10ul of the suggested working dilution to label 10 <sup>6</sup> or 100ul whole blood.  |
| <b>References</b>     | <ol style="list-style-type: none"><li>1. Hughes, G.J. <i>et al.</i> (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 <i>in vivo</i>. <a href="#">AIDS. 21: 1507-13.</a></li><li>2. Allen, J.S. <i>et al.</i> (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. <a href="#">Diabetes. 58: 138-45.</a></li><li>3. McIntosh, K. <i>et al.</i> (2006) The immunogenicity of human adipose-derived cells: temporal changes <i>in vitro</i>. <a href="#">Stem Cells. 24: 1246-53.</a></li><li>4. Sengstake, S. <i>et al.</i> (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. <a href="#">Int Immunol. 18 (7): 1171-8.</a></li><li>5. Villarreal Dorrego, M. <i>et al.</i> (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. <a href="#">Br J Dermatol. 154: 231-8.</a></li><li>6. Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. <a href="#">Blood. 118: 348-57.</a></li><li>7. Lacal, P.M. <i>et al.</i> (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. <a href="#">J Pharmacol Exp Ther. 347: 164-72.</a></li><li>8. Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. <a href="#">Blood. 118: 348-57.</a></li><li>9. Girbl, T. <i>et al.</i> (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. <a href="#">Cancer Res. 73: 561-70.</a></li><li>10. Hertzberg, L. <i>et al.</i> (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. <a href="#">Blood. 115: 1006-17.</a></li><li>11. Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. <a href="#">Diab Vasc Dis Res. 8: 221-8.</a></li><li>12. Dorvignit, D. <i>et al.</i> (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. <a href="#">MAbs. 4 (4): 488-96.</a></li><li>13. Karlsen, M. <i>et al.</i> (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. <a href="#">Scand J Immunol. 82 (6): 523-31.</a></li><li>14. Clark, L.E. <i>et al.</i> (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. <a href="#">Nat Commun. 9 (1): 1884.</a></li><li>15. Gu, Y. <i>et al.</i> (2019) Defining the structural basis for human alloantibody binding to human leukocyte antigen allele HLA-A*11:01. <a href="#">Nat Commun. 10 (1): 893.</a></li></ol> |
| <b>Storage</b>        | Store at +4°C or at -20°C if preferred.  |

---

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 18 months from date of despatch.

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 700 \(MCA928A700\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M352788:190408'

**Printed on 20 May 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)