

Datasheet: MCA1940APCT

Description:	MOUSE ANTI HUMAN CD19:APC
Specificity:	CD19
Format:	APC
Product Type:	Monoclonal Antibody
Clone:	LT19
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

RRID AB_1101049

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG conjugated to Allophycocyanin (APC) - lyophilised

Reconstitution Reconstitute with 0.25 ml distilled water

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	APC	650	661

Preparation Purified IgG from ascites prepared by precipitation and ion exchange chromatography

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide
Stabilisers 1% Bovine Serum Albumin

External Database Links

UniProt:
[P15391](#) [Related reagents](#)

Entrez Gene:
[930](#) CD19 [Related reagents](#)

Specificity	<p>Mouse anti Human CD19 antibody, clone LT19 recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells (de Rie et al. 1989).</p> <p>CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 (Bradbury et al. 1992), CD9, CD81 and CD82 (Horváth et al. 1998). CD19 is implicated in the down-regulation of B cell growth and proliferation (Pezzutto et al. 1987).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or cells or 100ul whole blood.
References	<ol style="list-style-type: none"> 1. Hughes, G.J. <i>et al.</i> (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 <i>in vivo</i>. AIDS. 21: 1507-13. 2. Allen, J.S. <i>et al.</i> (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. Diabetes. 58: 138-45. 3. McIntosh, K. <i>et al.</i> (2006) The immunogenicity of human adipose-derived cells: temporal changes <i>in vitro</i>. Stem Cells. 24: 1246-53. 4. Sengstake, S. <i>et al.</i> (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. Int Immunol. 18 (7): 1171-8. 5. Villarreal Dorrego, M. <i>et al.</i> (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. Br J Dermatol. 154: 231-8. 6. Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. Blood. 118: 348-57. 7. Lacal, P.M. <i>et al.</i> (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. J Pharmacol Exp Ther. 347: 164-72. 8. Franz, B. <i>et al.</i> (2011) <i>Ex vivo</i> characterization and isolation of rare memory B cells with antigen tetramers. Blood. 118: 348-57. 9. Girbl, T. <i>et al.</i> (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. Cancer Res. 73: 561-70. 10. Hertzberg, L. <i>et al.</i> (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. Blood. 115: 1006-17. 11. Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. Diab Vasc Dis Res. 8: 221-8. 12. Dorvignit, D. <i>et al.</i> (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. MAbs. 4 (4): 488-96. 13. Karlsen, M. <i>et al.</i> (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. Scand J Immunol. 82 (6): 523-31. 14. Clark, L.E. <i>et al.</i> (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. Nat Commun. 9 (1): 1884.
Storage	<p>Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected</p>

from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	12 months from date of reconstitution.
-------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:APC \(MCA928APC\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M343036:190110'

Printed on 10 Jan 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)