

Datasheet: MCA1940PE

Description:	MOUSE ANTI HUMAN CD19:RPE
Specificity:	CD19
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	LT19
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

RRID AB_323282

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution Reconstitute with 1 ml distilled water

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)

Stabilisers 1% Bovine Serum Albumin

5% Sucrose

External Database

Links

UniProt:

[P15391](#) [Related reagents](#)

Entrez Gene:

[930](#) CD19 [Related reagents](#)

Specificity **Mouse anti Human CD19 antibody, clone LT19** recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells ([de Rie et al. 1989](#)).

CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 ([Bradbury et al. 1992](#)), CD9, CD81 and CD82 ([Horváth et al. 1998](#)). CD19 is implicated in the down-regulation of B cell growth and proliferation ([Pezzutto et al. 1987](#)).

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells or cells or 100ul whole blood.

References

1. Hughes, G.J. *et al.* (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 *in vivo*. [AIDS. 21: 1507-13.](#)
2. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. [Diabetes. 58: 138-45.](#)
3. McIntosh, K. *et al.* (2006) The immunogenicity of human adipose-derived cells: temporal changes *in vitro*. [Stem Cells. 24: 1246-53.](#)
4. Sengstake, S. *et al.* (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. [Int Immunol. 18 \(7\): 1171-8.](#)
5. Villarreal Dorrego, M. *et al.* (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. [Br J Dermatol. 154: 231-8.](#)
6. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
7. Lacal, P.M. *et al.* (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. [J Pharmacol Exp Ther. 347: 164-72.](#)
8. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
9. Girbl, T. *et al.* (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. [Cancer Res. 73: 561-70.](#)
10. Hertzberg, L. *et al.* (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. [Blood. 115: 1006-17.](#)
11. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab Vasc Dis Res. 8: 221-8.](#)
12. Dorvignit, D. *et al.* (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. [MAbs. 4 \(4\): 488-96.](#)
13. Karlsen, M. *et al.* (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. [Scand J Immunol. 82 \(6\): 523-31.](#)
14. Clark, L.E. *et al.* (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. [Nat Commun. 9 \(1\): 1884.](#)
15. Gu, Y. *et al.* (2019) Defining the structural basis for human alloantibody binding to human leukocyte antigen allele HLA-A*11:01. [Nat Commun. 10 \(1\): 893.](#)

Storage Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of reconstitution.
------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M352801:190408'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)