

## Datasheet: MCA1940SBB580

<b>Description:</b>	MOUSE ANTI HUMAN CD19:StarBright Blue 580
<b>Specificity:</b>	CD19
<b>Format:</b>	StarBright Blue 580
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LT19
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/0.5ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human		
<b>Product Form</b>	Purified IgG conjugated to StarBright Blue 580 - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	StarBright Blue 580	475	582
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
<b>Buffer Solution</b>	Phosphate buffered saline		
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )		
	1% bovine serum albumin		
	0.1% Pluronic F68		
	0.1% PEG 3350		
	0.05% Tween 20		

UniProt:

[P15391](#)   [Related reagents](#)

Entrez Gene:

[930](#)   CD19   [Related reagents](#)

---

Specificity

**Mouse anti Human CD19 antibody, clone LT19** recognizes human CD19 also known as T-cell surface antigen Leu-12 or B-lymphocyte surface antigen B4. CD19 is a ~95 kDa type I single pass transmembrane glycoprotein expressed on follicular dendritic cells and B-cells during maturation but is lost on development into plasma cells ([de Rie et al. 1989](#)).

CD19 is the broadest lineage specific marker for B cells and functions as a B-cell co-receptor in conjunction with CD21 ([Bradbury et al. 1992](#)), CD9, CD81 and CD82 ([Horváth et al. 1998](#)). CD19 is implicated in the down-regulation of B cell growth and proliferation ([Pezzutto et al. 1987](#)).

---

Flow Cytometry

Use 5µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

---

References

1. Hughes, G.J. *et al.* (2007) Virus immunocapture provides evidence of CD8 lymphocyte-derived HIV-1 *in vivo*. [AIDS. 21: 1507-13.](#)
2. Allen, J.S. *et al.* (2009) Plasmacytoid dendritic cells are proportionally expanded at diagnosis of type 1 diabetes and enhance islet autoantigen presentation to T-cells through immune complex capture. [Diabetes. 58: 138-45.](#)
3. McIntosh, K. *et al.* (2006) The immunogenicity of human adipose-derived cells: temporal changes *in vitro*. [Stem Cells. 24: 1246-53.](#)
4. Sengstake, S. *et al.* (2006) CD21 and CD62L shedding are both inducible via P2X7Rs. [Int Immunol. 18 \(7\): 1171-8.](#)
5. Villarreal Dorrego, M. *et al.* (2006) Transfection of CD40 in a human oral squamous cell carcinoma keratinocyte line upregulates immune potency and costimulatory molecules. [Br J Dermatol. 154: 231-8.](#)
6. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
7. Lacal, P.M. *et al.* (2013) Glucocorticoid-induced tumor necrosis factor receptor family-related ligand triggering upregulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 and promotes leukocyte adhesion. [J Pharmacol Exp Ther. 347: 164-72.](#)
8. Franz, B. *et al.* (2011) *Ex vivo* characterization and isolation of rare memory B cells with antigen tetramers. [Blood. 118: 348-57.](#)
9. Girbl, T. *et al.* (2013) CD40-mediated activation of chronic lymphocytic leukemia cells promotes their CD44-dependent adhesion to hyaluronan and restricts CCL21-induced motility. [Cancer Res. 73: 561-70.](#)
10. Hertzberg, L. *et al.* (2010) Down syndrome acute lymphoblastic leukemia, a highly heterogeneous disease in which aberrant expression of CRLF2 is associated with mutated JAK2: a report from the International BFM Study Group. [Blood. 115: 1006-17.](#)
11. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab](#)

[Vasc Dis Res. 8: 221-8.](#)

12. Dorvignit, D. *et al.* (2012) Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: a case study. [MAbs. 4 \(4\): 488-96.](#)

13. Karlsen, M. *et al.* (2015) TLR-7 and -9 Stimulation of Peripheral Blood B Cells Indicate Altered TLR Signalling in Primary Sjögren's Syndrome Patients by Increased Secretion of Cytokines. [Scand J Immunol. 82 \(6\): 523-31.](#)

14. Clark, L.E. *et al.* (2018) Vaccine-elicited receptor-binding site antibodies neutralize two New World hemorrhagic fever arenaviruses. [Nat Commun. 9 \(1\): 1884.](#)

15. Gu, Y. *et al.* (2019) Defining the structural basis for human alloantibody binding to human leukocyte antigen allele HLA-A\*11:01. [Nat Commun. 10 \(1\): 893.](#)

16. Yang, C. *et al.* (2013) B cells promote tumor progression via STAT3 regulated-angiogenesis. [PLoS One. 8 \(5\): e64159.](#)

---

**Storage**

Store at +4°C.  
DO NOT FREEZE.  
This product should be stored undiluted.

---

**Guarantee**

12 months from date of despatch

---

**Acknowledgements**

This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

---

**Health And Safety Information**

Material Safety Datasheet documentation #20471 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA1940SBB580>  
20471

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M433251:241008'

**Printed on 10 Oct 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)