

Datasheet: MCA1949PET

Description:	MOUSE ANTI HUMAN CD29:RPE	
Specificity:	CD29	
Other names:	INTEGRIN BETA 1 CHAIN	
Format:	RPE	
Product Type:	Monoclonal Antibody	
Clone:	4B7R	
Isotype:	lgG1	
Quantity:	25 TESTS	

Product Details

RRID AB_1101774

Applications This product has been reported to work in the following applications. This information is derived

from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol

recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/2

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in

their own system using appropriate negative/positive controls.

Target Species Human **Product Form** Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution Reconstitute in 0.25 ml disilled water

Max Ex/Em **Fluorophore** Excitation Max (nm) **Emission Max (nm)** RPE 488nm laser 496 578

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide **Stabilisers**

Bovine Serum Albumin 1%

> 5% Sucrose

Occular melanoma cell line V+B2. **Immunogen**

External Database UniProt: Links

P05556 Related reagents

Entrez Gene:

3688 ITGB1 Related reagents

Synonyms

FNRB, MDF2, MSK12

Specificity

Mouse anti Human CD29 monoclonal antibody, clone 4B7R recognizes the human integrin beta 1 subunit, also known as CD29, a ~130 kDa (red) 115 kDa (non-red) single pass type I transmembrane glycoprotein expressed by most leucocytes and mesenchymal stem cells.

Integrin receptors are involved in the regulation of a variety of important biological functions, including embryonic development, wound repair, hemostasis and prevention of programmed cell death. They are also implicated in abnormal pathological states such as tumor directed angiogenesis, tumor cell growth, and metastasis. Surface expression of CD29 on human natural killer cells can be reduced by pretreatment with the glutathione-S-transferase inhibitor diethyl maleate (Horvath-Arcidiacono et al. 2003)

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Marshall, J.F. *et al.* (1998) Comparative analysis of integrins *in vitro* and *in vivo* in uveal and cutaneous melanomas. Br J Cancer. 77 (4): 522-9.
- 2. Pillay, J. *et al.* (2010) Functional heterogeneity and differential priming of circulating neutrophils in human experimental endotoxemia. J Leukoc Biol. 88: 211-20.
- 3. Kim, B.S. *et al.* (2011) Effects of the dichloromethane fraction of dipsaci radix on the osteoblastic differentiation of human alveolar bone marrow-derived mesenchymal stem cells. <u>Biosci Biotechnol</u> Biochem. 75:13-9.
- 4. Kato, H. *et al.* (2012) The primacy of β1 integrin activation in the metastatic cascade. <u>PLoS One.</u> 7 (10): e46576.
- 5. Meng, J. *et al.* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. <u>PLoS One. 6(3):e17454.</u>
- 6. Horvath-Arcidiacono, J.A. *et al.* (2003) Human natural killer cell activity against porcine targets: modulation by control of the oxidation-reduction environment and role of adhesion molecule interactions. Cell Immunol. 222: 35-44.
- 7. Tan, D.W. *et al.* (2013) Single-cell gene expression profiling reveals functional heterogeneity of undifferentiated human epidermal cells. <u>Development. 140 (7): 1433-44</u>.
- 8. Yang, J. *et al.* (2015) Cordycepin protected against the TNF-α-induced inhibition of osteogenic differentiation of human adipose-derived mesenchymal stem cells. <u>Int J Immunopathol Pharmacol.</u> 28 (3): 296-307.
- 9. Gu, Q. *et al.* (2015) *Ginkgo biloba* extract promotes osteogenic differentiation of human bone marrow mesenchymal stem cells in a pathway involving Wnt/β-catenin signaling. <u>Pharmacol Res.</u> 97: 70-8.
- 10. Chen, Y. *et al.* (2015) Effect of human umbilical cord mesenchymal stem cells transplantation on nerve fibers of a rat model of endometriosis. Int J Fertil Steril. 9 (1): 71-80.
- 11. Lee, H.J. *et al.* (2017) ICOSL expression in human bone marrow-derived mesenchymal stem cells promotes induction of regulatory T cells. <u>Sci Rep. 7: 44486.</u>
- 12. Yi, T. *et al.* (2015) Manufacture of Clinical-Grade Human Clonal Mesenchymal Stem Cell Products from Single Colony Forming Unit-Derived Colonies Based on the Subfractionation Culturing Method. <u>Tissue Eng Part C Methods</u>. 21 (12): 1251-62.
- 13. Sun, Y. *et al.* (2017) Antinociceptive Effect of Intrathecal Injection of Genetically Engineered Human Bone Marrow Stem Cells Expressing the Human Proenkephalin Gene in a Rat Model of Bone Cancer Pain. Pain Res Manag. 2017: 7346103.

Storage

Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Worldwide

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Tel: +49 (0) 89 8090 95 21

'M343042:190110'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint