

## Datasheet: MCA1949T

**BATCH NUMBER 160809**

<b>Description:</b>	MOUSE ANTI HUMAN CD29
<b>Specificity:</b>	CD29
<b>Other names:</b>	INTEGRIN BETA 1 CHAIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	4B7R
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting		▪		
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

**(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Ocular melanoma cell line V+B2.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P05556</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3688</a> ITGB1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	FNRB, MDF2, MSK12
<b>RRID</b>	AB_1101766
<b>Specificity</b>	<p><b>Mouse anti Human CD29 monoclonal antibody, clone 4B7R</b> recognizes the human integrin beta 1 subunit, also known as CD29, a ~130 kDa (red) 115 kDa (non-red) single pass type I transmembrane glycoprotein expressed by most leucocytes and <a href="#">mesenchymal stem cells</a>.</p> <p>Integrin receptors are involved in the regulation of a variety of important biological functions, including embryonic development, wound repair, hemostasis and prevention of programmed cell death. They are also implicated in abnormal pathological states such as tumor directed angiogenesis, tumor cell growth, and metastasis. Surface expression of CD29 on human natural killer cells can be reduced by pretreatment with the glutathione-S-transferase inhibitor diethyl maleate (<a href="#">Horvath-Arcidiacono et al. 2003</a>)</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Marshall, J.F. <i>et al.</i> (1998) Comparative analysis of integrins <i>in vitro</i> and <i>in vivo</i> in uveal and cutaneous melanomas. <a href="#">Br J Cancer. 77 (4): 522-9.</a></li> <li>2. Pillay, J. <i>et al.</i> (2010) Functional heterogeneity and differential priming of circulating neutrophils in human experimental endotoxemia. <a href="#">J Leukoc Biol. 88: 211-20.</a></li> <li>3. Kim, B.S. <i>et al.</i> (2011) Effects of the dichloromethane fraction of dipsaci radix on the osteoblastic differentiation of human alveolar bone marrow-derived mesenchymal stem cells. <a href="#">Biosci Biotechnol Biochem. 75:13-9.</a></li> <li>4. Kato, H. <i>et al.</i> (2012) The primacy of <math>\beta</math>1 integrin activation in the metastatic cascade. <a href="#">PLoS One. 7 (10): e46576.</a></li> <li>5. Meng, J. <i>et al.</i> (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. <a href="#">PLoS One. 6(3):e17454.</a></li> <li>6. Horvath-Arcidiacono, J.A. <i>et al.</i> (2003) Human natural killer cell activity against porcine</li> </ol>

targets: modulation by control of the oxidation-reduction environment and role of adhesion molecule interactions. [Cell Immunol. 222: 35-44.](#)

7. Tan, D.W. *et al.* (2013) Single-cell gene expression profiling reveals functional heterogeneity of undifferentiated human epidermal cells. [Development. 140 \(7\): 1433-44.](#)

8. Yang, J. *et al.* (2015) Cordycepin protected against the TNF- $\alpha$ -induced inhibition of osteogenic differentiation of human adipose-derived mesenchymal stem cells. [Int J Immunopathol Pharmacol. 28 \(3\): 296-307.](#)

9. Gu, Q. *et al.* (2015) *Ginkgo biloba* extract promotes osteogenic differentiation of human bone marrow mesenchymal stem cells in a pathway involving Wnt/ $\beta$ -catenin signaling. [Pharmacol Res. 97: 70-8.](#)

10. Chen, Y. *et al.* (2015) Effect of human umbilical cord mesenchymal stem cells transplantation on nerve fibers of a rat model of endometriosis. [Int J Fertil Steril. 9 \(1\): 71-80.](#)

11. Lee, H.J. *et al.* (2017) ICOSL expression in human bone marrow-derived mesenchymal stem cells promotes induction of regulatory T cells. [Sci Rep. 7: 44486.](#)

12. Yi, T. *et al.* (2015) Manufacture of Clinical-Grade Human Clonal Mesenchymal Stem Cell Products from Single Colony Forming Unit-Derived Colonies Based on the Subfractionation Culturing Method. [Tissue Eng Part C Methods. 21 \(12\): 1251-62.](#)

13. Sun, Y. *et al.* (2017) Antinociceptive Effect of Intrathecal Injection of Genetically Engineered Human Bone Marrow Stem Cells Expressing the Human Proenkephalin Gene in a Rat Model of Bone Cancer Pain. [Pain Res Manag. 2017: 7346103.](#)

14. Kim, S.H. *et al.* (2019) Forkhead box O1 (FOXO1) controls the migratory response of Toll-like receptor (TLR3)-stimulated human mesenchymal stromal cells. [J Biol Chem. 294 \(21\): 8424-37.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1949T>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- Goat Anti Mouse IgG (STAR77...) [HRP](#)
- Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
- Goat Anti Mouse IgG (STAR70...) [FITC](#)
- Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Goat Anti Mouse IgG (STAR76...)

[RPE](#)

Goat Anti Mouse IgG (H/L) (STAR117...)

[Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),  
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),  
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...)

[HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...)

[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...)

[FITC](#)

## Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M383123:210513'

Printed on 25 Mar 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)