

Datasheet: MCA1958FA

Description:	RAT ANTI MOUSE CD200:FITC
Specificity:	CD200
Other names:	OX2
Format:	FITC
<b>Product Type:</b>	Monoclonal Antibody
Clone:	OX-90
Isotype:	lgG2a
Quantity:	50 µg

## **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Immunogen	Mouse CD200-rat CD4 fusion protein.
External Database	UniProt:

O54901 Related reagents

**Entrez Gene:** 

17470 Cd200 Related reagents

 Synonyms
 Mox2

 RRID
 AB\_566612

Fusion Partners Spleen cells from immunised rats were fused with cells of the rat Y3 myeloma cell line

Rat anti Mouse CD200 antibody, clone OX-90 recognizes the mouse CD200 cell surface antigen, also known as OX2.

CD200 is expressed by splenic B lymphocytes, follicular dendritic cells, splenic endothelium and by neurons.

The CD200 - CD200 ligand system is of importance in the control of macrophage and granulocyte activation.

Flow Cytometry Use 10μl of the suggested working dilution to label 10<sup>6</sup> cells in 100μl.

The Fc region of monoclonal antibodies may bind to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR (<u>BUF041A/BUF041B</u>).

#### References

**Specificity** 

- 1. Hoek, R.M. *et al.* (2000) Down-regulation of the macrophage lineage through interaction with OX2 (CD200). <u>Science. 290 (5497): 1768-71.</u>
- 2. Nathan, C. & Muller, W.A. (2001) Putting the brakes on innate immunity: a regulatory role for CD200? Nat Immunol. 2 (1): 17-9.
- 3. Rijkers, E.S. (2007) Ligation of CD200R by CD200 is not required for normal murine myelopoiesis. <u>Eur J Haematol. 79: 410-6.</u>
- 4. Rijkers, E.S. *et al.* (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. <u>Mol Immunol. 45: 1126-35.</u>
- 5. Burger, P.E. *et al.* (2009) High aldehyde dehydrogenase activity: a novel functional marker of murine prostate stem/progenitor cells. <u>Stem Cells. 27: 2220-8.</u>
- 6. Ko, Y.C. *et al* (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. J Anat. 214: 183-95.
- 7. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. J Neuropathol Exp Neurol. 68: 159-67.
- 8. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. <u>Int Immunopharmacol. 11 (7): 794-801.</u>
- 9. Garza, L.A. *et al.* (2011) Bald scalp in men with androgenetic alopecia retains hair follicle stem cells but lacks CD200-rich and CD34-positive hair follicle progenitor cells. <u>J</u> Clin Invest. 121: 613-22.

- 10. Montiel, M. *et al.* (2015) Melatonin decreases brain apoptosis, oxidative stress, and CD200 expression and increased survival rate in mice infected by Venezuelan equine encephalitis virus. <u>Antivir Chem Chemother. 24 (3-4): 99-108.</u>
- 11. Liu, J.Q. *et al.* (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. <u>J Immunol. 197 (4): 1489-97.</u>
- 12. Liu, C. *et al.* (2018) The role of N-glycosylation of CD200-CD200R1 interaction in classical microglial activation. J Inflamm (Lond). 15: 28.
- 13. Tonecka, K. *et al.* (2021) The CD200 Regulates Inflammation in Mice Independently of TNF-α Production. Int J Mol Sci. 22 (10): 5358.
- 14. Pannunzio, B. *et al.* (2022) CD200R1 Contributes to Successful Functional Reinnervation after a Sciatic Nerve Injury Cells. 11 (11): 1786.

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1958FA">https://www.bio-rad-antibodies.com/SDS/MCA1958FA</a> 10041
Regulatory	For research purposes only

## Related Products

## **Recommended Negative Controls**

#### RAT IgG2a NEGATIVE CONTROL:FITC (MCA1212F)

Email: antibody\_sales\_us@bio-rad.com

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M412322'221110'

#### Printed on 19 Oct 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint