

## Datasheet: MCA1958T

**BATCH NUMBER 166282**

<b>Description:</b>	RAT ANTI MOUSE CD200
<b>Specificity:</b>	CD200
<b>Other names:</b>	OX2
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-90
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% sodium azide (NaN <sub>3</sub> )

<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0mg/ml
<b>Immunogen</b>	Mouse CD200-rat CD4 fusion protein.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">O54901</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">17470</a> Cd200    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Mox2
<b>RRID</b>	AB_2073825
<b>Fusion Partners</b>	Spleen cells from immunised rats were fused with cells of the rat Y3 myeloma cell line
<b>Specificity</b>	<p><b>Rat anti Mouse CD200 antibody, clone OX-90</b> recognizes the mouse CD200 cell surface antigen, also known as OX2.</p> <p>CD200 is expressed by splenic B lymphocytes, follicular dendritic cells, splenic endothelium and by neurons.</p> <p>The CD200 - CD200 ligand system is of importance in the control of macrophage and granulocyte activation.</p>
<b>Flow Cytometry</b>	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
<b>Histology Positive Control Tissue</b>	Mouse spleen
<b>References</b>	<ol style="list-style-type: none"> <li>1. Hoek, R.M. <i>et al.</i> (2000) Down-regulation of the macrophage lineage through interaction with OX2 (CD200). <a href="#">Science. 290 (5497): 1768-71.</a></li> <li>2. Nathan, C. &amp; Muller, W.A. (2001) Putting the brakes on innate immunity: a regulatory role for CD200? <a href="#">Nat Immunol. 2 (1): 17-9.</a></li> <li>3. Rijkers, E.S. (2007) Ligation of CD200R by CD200 is not required for normal murine myelopoiesis. <a href="#">Eur J Haematol. 79: 410-6.</a></li> <li>4. Rijkers, E.S. <i>et al.</i> (2008) The inhibitory CD200R is differentially expressed on human and mouse T and B lymphocytes. <a href="#">Mol Immunol. 45: 1126-35.</a></li> <li>5. Burger, P.E. <i>et al.</i> (2009) High aldehyde dehydrogenase activity: a novel functional marker of murine prostate stem/progenitor cells. <a href="#">Stem Cells. 27: 2220-8.</a></li> <li>6. Ko, Y.C. <i>et al.</i> (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. <a href="#">J Anat. 214: 183-95.</a></li> <li>7. Koning, N. <i>et al.</i> (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. <a href="#">J Neuropathol Exp Neurol. 68: 159-67.</a></li> </ol>

8. Seeds, R.E. *et al.* (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. [Int Immunopharmacol. 11 \(7\): 794-801.](#)
9. Garza, L.A. *et al.* (2011) Bald scalp in men with androgenetic alopecia retains hair follicle stem cells but lacks CD200-rich and CD34-positive hair follicle progenitor cells. [J Clin Invest. 121: 613-22.](#)
10. Montiel, M. *et al.* (2015) Melatonin decreases brain apoptosis, oxidative stress, and CD200 expression and increased survival rate in mice infected by Venezuelan equine encephalitis virus. [Antivir Chem Chemother. 24 \(3-4\): 99-108.](#)
11. Liu, J.Q. *et al.* (2016) A Critical Role for CD200R Signaling in Limiting the Growth and Metastasis of CD200+ Melanoma. [J Immunol. 197 \(4\): 1489-97.](#)
12. Liu, C. *et al.* (2018) The role of N-glycosylation of CD200-CD200R1 interaction in classical microglial activation. [J Inflamm \(Lond\). 15: 28.](#)
13. Lago, N. *et al.* (2018) CD200 modulates spinal cord injury neuroinflammation and outcome through CD200R1. [Brain Behav Immun. 73: 416-26.](#)
14. Tonecka, K. *et al.* (2021) The CD200 Regulates Inflammation in Mice Independently of TNF- $\alpha$  Production. [Int J Mol Sci. 22 \(10\): 5358.](#)
15. Pannunzio, B. *et al.* (2022) CD200R1 Contributes to Successful Functional Reinnervation after a Sciatic Nerve Injury [Cells. 11 \(11\): 1786.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1958T>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight@800</a>
Rabbit Anti Rat IgG (STAR17...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Goat Anti Rat IgG (STAR69...)	<a href="#">FITC</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@800</a>
Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>

### Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M412318:221110'

**Printed on 20 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)