

Datasheet: MCA1960F BATCH NUMBER 161742

Description:	MOUSE ANTI HUMAN CD200:FITC	
Specificity:	CD200	
Other names:	OX2	
Format:	FITC	
Product Type:	Monoclonal Antibody	
Clone:	OX-104	
Isotype:	lgG1	
Quantity:	0.1 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			Neat - 1/5

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

roduct Form	Purified IgG conjugate	ad to Fluorescein leath	niocvanate Isomer :
	r united tyo conjugate	ed to i idorescent isoti	nocyanate isomei
ax Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	FITC	490	525
reparation	Purified InG prepared	by affinity chromatog	ranhy on Protein A
paration fer Solution	Purified IgG prepared supernatant Phosphate buffered s		raphy on Protein A

Concentrations

External	Database
Links	

UniProt:

P41217 Related reagents

Entrez Gene:

4345 CD200 Related reagents

Synonyms

MOX1, MOX2

RRID

AB_323322

Specificity

Mouse anti Human CD200 antibody, clone OX-104 recognizes the human CD200 cell surface antigen, also known as OX2.

CD200 is expressed by a subset of B lymphocytes, some endothelial cells and by neurons. The CD200-CD200 ligand system is of importance in the control of macrophage and granulocyte activation.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Wright, G.J. *et al.* (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. <u>Immunology 102 (2): 173-9.</u>
- 2. Ko, Y.C. *et al.* (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. <u>J Anat. 214: 183-95.</u>
- 3. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. <u>J Neuropathol Exp Neurol</u>. 68: 159-67.
- 4. Koning, N. *et al.* (2007) Downregulation of macrophage inhibitory molecules in multiple sclerosis lesions. <u>Ann Neurol. 62: 504-14.</u>
- 5. Raftery, M.J. *et al.* (2004) Shaping phenotype, function, and survival of dendritic cells by cytomegalovirus-encoded IL-10. <u>J Immunol. 173: 3383-91.</u>
- 6. Meuth, S.G. *et al.* (2008) CNS inflammation and neuronal degeneration is aggravated by impaired CD200-CD200R-mediated macrophage silencing. <u>J Neuroimmunol. 194:</u> 62-9.
- 7. Yamauchi, K. and Kurosaka, A. (2010) Expression and function of glycogen synthase kinase-3 in human hair follicles. <u>Arch Dermatol Res. 302: 263-70.</u>
- 8. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful in situ markers for human epithelial hair follicle stem cells and their niche. Exp Dermatol. 17: 592-609.
- 9. Ohyama, M. *et al.* (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. <u>J Clin Invest. 116: 249-60.</u>
- 10. Darmochwal-Kolarz, D. *et al.* (2013) The expressions of co-stimulatory molecules are altered on putative antigen-presenting cells in cord blood. <u>Am J Reprod Immunol. 69 (2):</u> 180-7.
- 11. Colmont, C.S. *et al.* (2013) CD200-expressing human basal cell carcinoma cells initiate tumor growth. Proc Natl Acad Sci U S A. 110 (4): 1434-9.

- 12. Chen, H.J. *et al.* (2015) Human placenta-derived adherent cells improve cardiac performance in mice with chronic heart failure. <u>Stem Cells Transl Med. 4 (3): 269-75.</u>
- 13. Ohyama, M. & Kobayashi, T. (2012) Isolation and characterization of stem cell-enriched human and canine hair follicle keratinocytes. <u>Methods Mol Biol. 879:</u> 389-401.
- 14. Patel, G.K. *et al.* (2012) Identification and characterization of tumor-initiating cells in human primary cutaneous squamous cell carcinoma. <u>J Invest Dermatol. 132 (2): 401-9.</u>
 15. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful *in situ* markers for human epithelial hair follicle stem cells and their niche. Exp Dermatol. 17 (7): 592-609.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1960F 10041
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Email: antibody_sales_us@bio-rad.com

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385246:210513'

Printed on 18 Jan 2024