

# Datasheet: MCA1960FT

**BATCH NUMBER 165927**

<b>Description:</b>	MOUSE ANTI HUMAN CD200:FITC
<b>Specificity:</b>	CD200
<b>Other names:</b>	OX2
<b>Format:</b>	FITC
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-104
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/5

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% sodium azide (NaN <sub>3</sub> )		
Stabilisers	1% bovine serum albumin		
Approx. Protein	IgG concentration 0.1mg/ml		

## Concentrations

### External Database

#### Links

#### UniProt:

[P41217](#)

[Related reagents](#)

#### Entrez Gene:

[4345](#)

CD200

[Related reagents](#)

#### Synonyms

MOX1, MOX2

#### RRID

AB\_2074068

#### Specificity

**Mouse anti Human CD200 antibody, clone OX-104** recognizes the human CD200 cell surface antigen, also known as OX2.

CD200 is expressed by a subset of B lymphocytes, some endothelial cells and by neurons. The CD200-CD200 ligand system is of importance in the control of macrophage and granulocyte activation.

#### Flow Cytometry

Use 10µl of the suggested working dilution to label 10<sup>6</sup> cells in 100µl

#### References

1. Wright, G.J. *et al.* (2001) The unusual distribution of the neuronal/lymphoid cell surface CD200 (OX2) glycoprotein is conserved in humans. [Immunology 102 \(2\): 173-9.](#)
2. Raftery, M.J. *et al.* (2004) Shaping phenotype, function, and survival of dendritic cells by cytomegalovirus-encoded IL-10. [J Immunol. 173: 3383-91.](#)
3. Ohyama, M. *et al.* (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. [J Clin Invest. 116: 249-60.](#)
4. Koning, N. *et al.* (2007) Downregulation of macrophage inhibitory molecules in multiple sclerosis lesions. [Ann Neurol. 62: 504-14.](#)
5. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful *in situ* markers for human epithelial hair follicle stem cells and their niche. [Exp Dermatol. 17 \(7\): 592-609.](#)
6. Kloepper, J.E. *et al.* (2008) Immunophenotyping of the human bulge region: the quest to define useful *in situ* markers for human epithelial hair follicle stem cells and their niche. [Exp Dermatol. 17: 592-609.](#)
7. Meuth, S.G. *et al.* (2008) CNS inflammation and neuronal degeneration is aggravated by impaired CD200-CD200R-mediated macrophage silencing. [J Neuroimmunol. 194: 62-9.](#)
8. Koning, N. *et al.* (2009) Distribution of the immune inhibitory molecules CD200 and CD200R in the normal central nervous system and multiple sclerosis lesions suggests neuron-glia and glia-glia interactions. [J Neuropathol Exp Neurol. 68: 159-67.](#)
9. Ko, Y.C. *et al.* (2009) Endothelial CD200 is heterogeneously distributed, regulated and involved in immune cell-endothelium interactions. [J Anat. 214: 183-95.](#)
10. Yamauchi, K. and Kurosaka, A. (2010) Expression and function of glycogen synthase kinase-3 in human hair follicles. [Arch Dermatol Res. 302: 263-70.](#)
11. Patel, G.K. *et al.* (2012) Identification and characterization of tumor-initiating cells in human primary cutaneous squamous cell carcinoma. [J Invest Dermatol. 132 \(2\): 401-9.](#)

12. Ohyama, M. & Kobayashi, T. (2012) Isolation and characterization of stem cell-enriched human and canine hair follicle keratinocytes. [Methods Mol Biol. 879: 389-401.](#)
13. Colmont, C.S. *et al.* (2013) CD200-expressing human basal cell carcinoma cells initiate tumor growth. [Proc Natl Acad Sci U S A. 110 \(4\): 1434-9.](#)
14. Darmochwal-Kolarz, D. *et al.* (2013) The expressions of co-stimulatory molecules are altered on putative antigen-presenting cells in cord blood. [Am J Reprod Immunol. 69 \(2\): 180-7.](#)
15. Chen, H.J. *et al.* (2015) Human placenta-derived adherent cells improve cardiac performance in mice with chronic heart failure. [Stem Cells Transl Med. 4 \(3\): 269-75.](#)
16. Bertolini, M. *et al.* (2023) Mechanical epilation exerts complex biological effects on human hair follicles and perifollicular skin: An *ex vivo* study approach. [Int J Cosmet Sci. Nov 03 \[Epub ahead of print\].](#)

<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1960FT10041">https://www.bio-rad-antibodies.com/SDS/MCA1960FT10041</a></p>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
 'M419193:230606'

Printed on 18 Jan 2024