

Datasheet: MCA1965GA

BATCH NUMBER 1804

| | |
|----------------------|--------------------------------|
| Description: | MOUSE ANTI HUMAN CD91 |
| Specificity: | CD91 |
| Other names: | ALPHA-2 MACROGLOBULIN RECEPTOR |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | A2Mr alpha-2 |
| Isotype: | IgG1 |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|-------------------------|
| Flow Cytometry | ▪ | | | 1/50 1/100 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | ▪ | | | Direct |
| Immunoprecipitation | ▪ | | | |
| Western Blotting | ▪ | | | Non-reducing conditions |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------|---|
| Target Species | Human |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |

| | |
|-----------------------------------|--|
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml |
| Immunogen | Purified alpha2 macroglobulin receptor |
| External Database Links | <p>UniProt: Q07954 Related reagents</p> <p>Entrez Gene: 4035 LRP1 Related reagents</p> |
| Synonyms | A2MR, APR |
| RRID | AB_323544 |
| Fusion Partners | Spleen cells from immunised Balb/c mice were fused with cells of the NS1 mouse myeloma cell line. |
| Specificity | <p>Mouse anti Human CD91 antibody, clone A2Mr alpha-2 recognizes human CD91, also known as Prolow-density lipoprotein receptor-related protein 1, Alpha-2-macroglobulin receptor or apolipoprotein E receptor. CD91 is a 4525 amino acid protein post translationally cleaved into 3 subunits, a 85 kDa type I transmembrane carboxyl chain (LRP85) non-covalently bound to a 515 kDa extracellular N-terminal subunit (LRP515)containing multiple EGF-like and LDL-receptor Class A and Class B domains. Additionally, there is an intracellular domain (LRPICD) which can be cleaved from the transmambrane domain by gamma secretase (May et al. 2004). Clone A2Mr alpha-2 detects an epitope within the LRP515 chain.</p> <p>CD91 is a multifunctional protein involved in processes including the phagocytosis and endocytosis of apoptotic cells (Nilsson et al. 2012), clearance of activated serum alpha-2-macroglobulin (Kristensen et al. 1990), modulation of the inflammatory response (Staudt et al. 2013) and acts as a receptor for <i>Pseudomonas aeruginosa</i> exotoxin A (Kounnas et al. 1992).</p> <p>Mouse anti Human CD91, clone A2Mr alpha-2 has been used extensively for the detection of CD91 by flow cytometry and immunohistochemistry on formalin fixed paraffin embedded tissues (Bourazopoulou et al. 2009).</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
| Histology Positive Control Tissue | Liver |
| References | 1. Moestrup, S.K. <i>et al.</i> (1992) Distribution of the alpha 2-macroglobulin receptor/low density lipoprotein receptor-related protein in human tissues. Cell Tissue Res. 269 (3): 375-82. |

2. Moestrup, S.K. & Hokland, P. (1992) Surface expression of the alpha 2-macroglobulin receptor on human malignant blood cells. [Leuk Res. 16 \(3\): 227-34.](#)
3. Chen, J.S. *et al.* (2010) Secreted Heat Shock Protein 90{alpha} Induces Colorectal Cancer Cell Invasion through CD91/LRP-1 and NF-{kappa}B-mediated Integrin {alpha}V Expression. [J Biol Chem. 285: 25458-66.](#)
4. Bourazopoulou, E. *et al.* (2009) Functional expression of the alpha 2-macroglobulin receptor CD91 in salivary gland epithelial cells. [J Autoimmun. 33: 141-6.](#)
5. Hvidberg, V. *et al.* (2005) Identification of the receptor scavenging hemopexin-heme complexes. [Blood. 106: 2572-9.](#)
6. Hodge, S. *et al.* (2006) Azithromycin increases phagocytosis of apoptotic bronchial epithelial cells by alveolar macrophages. [Eur Respir J. 28 \(3\): 486-95.](#)
7. Hodge, S. *et al.* (2007) Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. [Am J Respir Cell Mol Biol. 37: 748-55.](#)
8. Pawluczyk, I.Z. *et al.* (2008) Perindoprilat modulates the activity of lipoprotein receptor-related protein in human mesangial cells. [J Biol Chem. 283: 4588-94.](#)
9. Cunningham, A.J. *et al.* (2012) Prolonged neutrophil dysfunction after *Plasmodium falciparum* malaria is related to hemolysis and heme oxygenase-1 induction. [J Immunol. 189 \(11\): 5336-46.](#)
10. Inage, E. *et al.* (2014) Critical Roles for PU.1, GATA1, and GATA2 in the expression of human FcεRI on mast cells: PU.1 and GATA1 transactivate FCER1A, and GATA2 transactivates FCER1A and MS4A2. [J Immunol. 192 \(8\): 3936-46.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1965GA>
10040

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

| | |
|---|----------------------|
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | HRP |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Goat Anti Mouse IgG (STAR70...) | FITC |

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M366018:200529'

Printed on 19 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)