

Datasheet: MCA1965GA

Description:	MOUSE ANTI HUMAN CD91
Specificity:	CD91
Other names:	ALPHA-2 MACROGLOBULIN RECEPTOR
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	A2Mr alpha-2
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			Direct
Immunoprecipitation	▪			
Western Blotting	▪			Non-reducing conditions

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Purified alpha2 macroglobulin receptor
External Database Links	<p>UniProt: Q07954 Related reagents</p> <p>Entrez Gene: 4035 LRP1 Related reagents</p>
Synonyms	A2MR, APR
RRID	AB_323544
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	<p>Mouse anti Human CD91 antibody, clone A2Mr alpha-2 recognizes human CD91, also known as Prolow-density lipoprotein receptor-related protein 1, Alpha-2-macroglobulin receptor or apolipoprotein E receptor. CD91 is a 4525 amino acid protein post translationally cleaved into 3 subunits, a 85 kDa type I transmembrane carboxyl chain (LRP85) non-covalently bound to a 515 kDa extracellular N-terminal subunit (LRP515)containing multiple EGF-like and LDL-receptor Class A and Class B domains. Additionally, there is an intracellular domain (LRPICD) which can be cleaved from the transmambrane domain by gamma secretase (May et al. 2004). Clone A2Mr alpha-2 detects an epitope within the LRP515 chain.</p> <p>CD91 is a multifunctional protein involved in processes including the phagocytosis and endocytosis of apoptotic cells (Nilsson et al. 2012), clearance of activated serum alpha-2-macroglobulin (Kristensen et al. 1990), modulation of the inflammatory response (Staudt et al. 2013) and acts as a receptor for <i>Pseudomonas aeruginosa</i> exotoxin A (Kounnas et al. 1992).</p> <p>Mouse anti Human CD91, clone A2Mr alpha-2 has been used extensively for the detection of CD91 by flow cytometry and immunohistochemistry on formalin fixed paraffin embedded tissues (Bourazopoulou et al. 2009).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
Histology Positive Control Tissue	Human liver
References	1. Moestrup, S.K. <i>et al.</i> (1992) Distribution of the alpha 2-macroglobulin receptor/low density lipoprotein receptor-related protein in human tissues. Cell Tissue Res. 269 (3): 375-82.

2. Moestrup, S.K. & Hokland, P. (1992) Surface expression of the alpha 2-macroglobulin receptor on human malignant blood cells. [Leuk Res. 16 \(3\): 227-34.](#)
3. Hvidberg, V. *et al.* (2005) Identification of the receptor scavenging hemopexin-heme complexes. [Blood. 106: 2572-9.](#)
4. Hodge, S. *et al.* (2006) Azithromycin increases phagocytosis of apoptotic bronchial epithelial cells by alveolar macrophages. [Eur Respir J. 28 \(3\): 486-95.](#)
5. Hodge, S. *et al.* (2007) Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. [Am J Respir Cell Mol Biol. 37: 748-55.](#)
6. Pawluczyk, I.Z. *et al.* (2008) Perindoprilat modulates the activity of lipoprotein receptor-related protein in human mesangial cells. [J Biol Chem. 283: 4588-94.](#)
7. Bourazopoulou, E. *et al.* (2009) Functional expression of the alpha 2-macroglobulin receptor CD91 in salivary gland epithelial cells. [J Autoimmun. 33: 141-6.](#)
8. Chen, J.S. *et al.* (2010) Secreted Heat Shock Protein 90{alpha} Induces Colorectal Cancer Cell Invasion through CD91/LRP-1 and NF- κ B-mediated Integrin {alpha}V Expression. [J Biol Chem. 285: 25458-66.](#)
9. Cunnington, A.J. *et al.* (2012) Prolonged neutrophil dysfunction after *Plasmodium falciparum* malaria is related to hemolysis and heme oxygenase-1 induction. [J Immunol. 189 \(11\): 5336-46.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1965GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC

Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M412374:221111'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)