

Datasheet: MCA1965PE

BATCH NUMBER 150607

| | |
|----------------------|--------------------------------|
| Description: | MOUSE ANTI HUMAN CD91:RPE |
| Specificity: | CD91 |
| Other names: | ALPHA-2 MACROGLOBULIN RECEPTOR |
| Format: | RPE |
| Product Type: | Monoclonal Antibody |
| Clone: | A2Mr alpha-2 |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | | | |
|------------------------|---|----------------------------|--------------------------|
| Target Species | Human | | |
| Product Form | Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized | | |
| Reconstitution | Reconstitute with 1.0 ml distilled water | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | RPE 488nm laser | 496 | 578 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative | 0.09% Sodium Azide | | |
| Stabilisers | 1% Bovine Serum Albumin | | |

5% Sucrose

| | |
|------------------|--|
| Immunogen | Purified alpha2 macroglobulin receptor |
|------------------|--|

| | |
|--------------------------------|---|
| External Database Links | UniProt: Q07954 Related reagents |
| | Entrez Gene: 4035 LRP1 Related reagents |

| | |
|-----------------|-----------|
| Synonyms | A2MR, APR |
|-----------------|-----------|

| | |
|-------------|-----------|
| RRID | AB_323268 |
|-------------|-----------|

| | |
|------------------------|---|
| Fusion Partners | Spleen cells from immunised Balb/c mice were fused with cells of the NS1 mouse myeloma cell line. |
|------------------------|---|

| | |
|--------------------|---|
| Specificity | <p>Mouse anti Human CD91 antibody, clone A2Mr alpha-2 recognizes human CD91, also known as Prolow-density lipoprotein receptor-related protein 1, Alpha-2-macroglobulin receptor or apolipoprotein E receptor. CD91 is a 4525 amino acid protein post translationally cleaved into 3 subunits, a 85 kDa type I transmembrane carboxyl chain (LRP85) non-covalently bound to a 515 kDa extracellular N-terminal subunit (LRP515) containing multiple EGF-like and LDL-receptor Class A and Class B domains. Additionally, there is an intracellular domain (LRPICD) which can be cleaved from the transmembrane domain by gamma secretase (May et al. 2004). Clone A2Mr alpha-2 detects an epitope within the LRP515 chain.</p> <p>CD91 is a multifunctional protein involved in processes including the phagocytosis and endocytosis of apoptotic cells (Nilsson et al. 2012), clearance of activated serum alpha-2-macroglobulin (Kristensen et al. 1990), modulation of the inflammatory response (Staudt et al. 2013) and acts as a receptor for <i>Pseudomonas aeruginosa</i> exotoxin A (Kounnas et al. 1992).</p> <p>Mouse anti Human CD91, clone A2Mr alpha-2 has been used extensively for the detection of CD91 by flow cytometry and immunohistochemistry on formalin fixed paraffin embedded tissues (Bourazopoulou et al. 2009).</p> |
|--------------------|---|

| | |
|-----------------------|---|
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul. |
|-----------------------|---|

| | |
|-------------------|---|
| References | <ol style="list-style-type: none">1. Moestrup, S.K. <i>et al.</i> (1992) Distribution of the alpha 2-macroglobulin receptor/low density lipoprotein receptor-related protein in human tissues. Cell Tissue Res. 269 (3): 375-82.2. Moestrup, S.K. & Hokland, P. (1992) Surface expression of the alpha 2-macroglobulin receptor on human malignant blood cells. Leuk Res. 16 (3): 227-34.3. Chen, J.S. <i>et al.</i> (2010) Secreted Heat Shock Protein 90{alpha} Induces Colorectal Cancer Cell Invasion through CD91/LRP-1 and NF-κB-mediated Integrin {alpha}V Expression. J Biol Chem. 285: 25458-66. |
|-------------------|---|

4. Bourazopoulou, E. *et al.* (2009) Functional expression of the alpha 2-macroglobulin receptor CD91 in salivary gland epithelial cells. [J Autoimmun. 33: 141-6.](#)
5. Hvidberg, V. *et al.* (2005) Identification of the receptor scavenging hemopexin-heme complexes. [Blood. 106: 2572-9.](#)
6. Hodge, S. *et al.* (2006) Azithromycin increases phagocytosis of apoptotic bronchial epithelial cells by alveolar macrophages. [Eur Respir J. 28 \(3\): 486-95.](#)
7. Hodge, S. *et al.* (2007) Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. [Am J Respir Cell Mol Biol. 37: 748-55.](#)
8. Pawluczyk, I.Z. *et al.* (2008) Perindoprilat modulates the activity of lipoprotein receptor-related protein in human mesangial cells. [J Biol Chem. 283: 4588-94.](#)
9. Cunnington, A.J. *et al.* (2012) Prolonged neutrophil dysfunction after *Plasmodium falciparum* malaria is related to hemolysis and heme oxygenase-1 induction. [J Immunol. 189 \(11\): 5336-46.](#)
10. Inage, E. *et al.* (2014) Critical Roles for PU.1, GATA1, and GATA2 in the expression of human FcεRI on mast cells: PU.1 and GATA1 transactivate FCER1A, and GATA2 transactivates FCER1A and MS4A2. [J Immunol. 192 \(8\): 3936-46.](#)

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1965PE>
20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M375399:210104'

