Datasheet: MCA1965PET

Description:	MOUSE ANTI HUMAN CD91:RPE
Specificity:	CD91
Other names:	ALPHA-2 MACROGLOBULIN RECEPTOR
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	A2Mr alpha-2
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution

Reconstitute in 0.25 ml disilled water

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09%	Sodium Azide
1%	Bovine Serum Albumin
5%	Sucrose

Immunogen	Purified alpha2 macroglobulin receptor
External Database Links	**UniProt:** Q07954 Related reagents **Entrez Gene:** 4035 LRP1 Related reagents
Synonyms	A2MR, APR
RRID	AB_1102457
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the NS1 mouse myeloma cell line.
Specificity	**Mouse anti Human CD91 antibody, clone A2Mr alpha-2** recognizes human CD91, also known as Prolow-density lipoprotein receptor-related protein 1, Alpha-2-macroglobulin receptor or apolipoprotein E receptor. CD91 is a 4525 amino acid protein post translationally cleaved into 3 subunits, a 85 kDa type I transmembrane carboxyl chain (LRP85) non-covalently bound to a 515 kDa extracellular N-terminal subunit (LRP515)containing multiple EGF-like and LDL-receptor Class A and Class B domains. Additionally, there is an intracellular domain (LRPICD) which can be cleaved from the transmambrane domain by gamma secretase (May *et al.* 2004). Clone A2Mr alpha-2 detects an epitope within the LRP515 chain. CD91 is a multifunctional protein involved in processes inluding the phagocytosis and endocytosis of apoptotic cells (Nilsson *et al.* 2012), clearance of activated serum alpha-2-macroglobulin (Kristensen *et al.* 1990), modulation of the inflammatory response (Staudt *et al.* 2013) and acts as a receptor for *Pseudomonas aeruginosa* exotoxin A (Kounnas *et al.* 1992). Mouse anti Human CD91, clone A2Mr alpha-2 has been used extensively for the detection of CD91 by flow cytometry and immunohistochemistry on formalin fixed paraffin embedded tissues (Bourazopoulou *et al.* 2009).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.
References	1. Moestrup, S.K. *et al.* (1992) Distribution of the alpha 2-macroglobulin receptor/low density lipoprotein receptor-related protein in human tissues. Cell Tissue Res. 269 (3): 375-82. 2. Moestrup, S.K. & Hokland, P. (1992) Surface expression of the alpha 2-macroglobulin receptor on human malignant blood cells. Leuk Res. 16 (3): 227-34. 3. Chen, J.S. *et al.* (2010) Secreted Heat Shock Protein 90{alpha} Induces Colorectal Cancer Cell Invasion through CD91/LRP-1 and NF-{kappa}B-mediated Integrin {alpha}V Expression. J Biol Chem. 285: 25458-66. 4. Bourazopoulou, E. *et al.* (2009) Functional expression of the alpha 2-macroglobulin receptor CD91 in salivary gland epithelial cells. J Autoimmun. 33: 141-6.

5. Hvidberg, V. *et al.* (2005) Identification of the receptor scavenging hemopexin-heme complexes. Blood. 106: 2572-9.
6. Hodge, S. *et al.* (2006) Azithromycin increases phagocytosis of apoptotic bronchial epithelial cells by alveolar macrophages. Eur Respir J. 28 (3): 486-95.
7. Hodge, S. *et al.* (2007) Smoking alters alveolar macrophage recognition and phagocytic ability: implications in chronic obstructive pulmonary disease. Am J Respir Cell Mol Biol. 37: 748-55.
8. Pawluczyk, I.Z. *et al.* (2008) Perindoprilat modulates the activity of lipoprotein receptor-related protein in human mesangial cells. J Biol Chem. 283: 4588-94.
9. Cunnington, A.J. *et al.* (2012) Prolonged neutrophil dysfunction after *Plasmodium falciparum* malaria is related to hemolysis and heme oxygenase-1 induction. J Immunol. 189 (11): 5336-46.
10. Inage, E. *et al.* (2014) Critical Roles for PU.1, GATA1, and GATA2 in the expression of human FcεRI on mast cells: PU.1 and GATA1 transactivate FCER1A, and GATA2 transactivates FCER1A and MS4A2. J Immunol. 192 (8): 3936-46.

Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A)
HUMAN SEROBLOCK (BUF070B)

North & South America Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com
Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com
Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Printed on 09 Feb 2021