# Datasheet: MCA1967 BATCH NUMBER 167027

Description:	RAT ANTI MOUSE CD44v6
Specificity:	CD44v6
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	9A4
lsotype:	lgG1
Quantity:	0.1 mg

#### **Product Details**

Applications	This product has been reported to work in the following applications. This information is
	derived from testing within our laboratories, peer-reviewed publications or personal
	communications from the originators. Please refer to references indicated for further
	information. For general protocol recommendations, please visit <u>www.bio-</u>
	rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen				
Immunohistology - Paraffin (1)				
ELISA	•			
Immunoprecipitation			-	
Western Blotting	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections.Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Antibody purified from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% sodium azide (NaN <sub>3</sub> )

Approx. Protein Concentrations	IgG concentration 0.5mg/ml
Immunogen	GST-CD44v6 fusion protein.
External Database Links	UniProt:         P15379       Related reagents         Entrez Gene:         12505       Cd44       Related reagents
Synonyms	Ly-24
RRID	AB_323213
Fusion Partners	Spleen cells from immunized female DA rats were fused with cells from the SP/0 myeloma.
Specificity	Rat anti Mouse CD44v6, clone 9A4 recognizes an epitope encoded by exon v6 on the variant portion of murine CD44. CD44v6 has been identified along with ICAM-1 as a co-receptor for Met and in mammary epithelia appears to act at the primary Met co-receptor (Di-Cicco <i>et al.</i> 2015). Rat anti Mouse CD44v6, clone 9A4 has been used successfully for the identification of CD44v6 expressing cells by immunofluorescence (Di-Cicco <i>et al.</i> 2015), western blotting (Lo <i>et al.</i> 2012). Immunohistochemistry (Weilenga <i>et al.</i> 1999) along with ELISA and flow cytometry (Weiss <i>et al.</i> 1997).
References	<ol> <li>Weiss JM <i>et al.</i> (1997) An essential role for CD44 variant isoforms in epidermal Langerhans cell and blood dendritic cell function. J Cell Biol. 137 (5): 1137-47.</li> <li>Wielenga VJ <i>et al.</i> (1999) Expression of CD44 in Apc and Tcf mutant mice implies regulation by the WNT pathway. Am J Pathol. 154 (2): 515-23.</li> <li>Katagiri, Y.U. <i>et al.</i> (1999) CD44 variants but not CD44s cooperate with beta1- containing integrins to permit cells to bind to osteopontin independently of arginine- glycine-aspartic acid, thereby stimulating cell motility and chemotaxis. Cancer Res. 59: 219-26.</li> <li>Hebbard, L. <i>et al.</i> (2000) CD44 expression and regulation during mammary gland development and function. J Cell Sci. 113: 2619-30.</li> <li>Rochman, M. <i>et al.</i> (2000) The CD44 receptor of lymphoma cells: structure-function relationships and mechanism of activation. Cell Adhes Commun. 7: 331-47.</li> <li>Khaldoyanidi, S. <i>et al.</i> (2002) CD44 variant-specific antibodies trigger hemopoiesis by selective release of cytokines from bone marrow macrophages. Blood. 99: 3955-61.</li> <li>Termeer, C. <i>et al.</i> (2003) Targeting dendritic cells with CD44 monoclonal antibodies selectively inhibits the proliferation of naive CD4+ T-helper cells by induction of FAS-independent T-cell apoptosis. Immunology. 109: 32-40.</li> <li>Matrosova, V.Y. <i>et al.</i> (2004) Hyaluronic acid facilitates the recovery of hematopoiesis following 5-fluorouracil administration. Stem Cells. 22: 544-55.</li> </ol>

	9. Serobyan, N. et al. (2005) Nicolinic acetylcholine receptor-mediated sumulation of				
	endothelial cells results in the arrest of haematopoietic progenitor cells on endothelium. Br				
	<u>J Haematol. 129: 257-65.</u>				
	10. Boon, E.M. et al. (2006) Activation of Wnt signaling in the intestinal mucosa of Apc				
	+/min mice does not cause overexpression of the receptor tyrosine kinase Met. Cancer				
	<u>Sci. 97: 710-5.</u>				
	11. Wallach-Dayan, S.B. et al. (2008) DNA vaccination with CD44 variant isoform reduces				
	mammary tumor local growth and lung metastasis. Mol Cancer Ther. 7: 1615-23.				
	12. Lo, B. et al. (2012) Lkb1 regulates organogenesis and early oncogenesis along				
	AMPK-dependent and -independent pathways. J Cell Biol. 199: 1117-30.				
	13. Nakagawa, H. et al. (2014) Loss of liver E-cadherin induces sclerosing cholangitis and				
	promotes carcinogenesis. <u>Proc Natl Acad Sci U S A. 111(3):1090-5.</u>				
	14. Di-Cicco, A. et al. (2015) Paracrine Met signaling triggers epithelial-mesenchymal				
	transition in mammary luminal progenitors, affecting their fate. Elife. 2015 Jul 13;4.				
	15. Shibata, W. et al. (2017) Helicobacter-induced gastric inflammation alters the				
	properties of gastric tissue stem/progenitor cells. BMC Gastroenterol. 17 (1): 145.				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.				
Storage Guarantee	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.				
Storage Guarantee Health And Safety	<ul> <li>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</li> <li>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</li> <li>12 months from date of despatch</li> </ul>				
Storage Guarantee Health And Safety Information	<ul> <li>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</li> <li>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</li> <li>12 months from date of despatch</li> <li>Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1967</li> </ul>				
Storage Guarantee Health And Safety Information	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1967 10040				
Storage Guarantee Health And Safety Information	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.         Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.         12 months from date of despatch         Material Safety Datasheet documentation #10040 available at:         https://www.bio-rad-antibodies.com/SDS/MCA1967         10040				
Storage Guarantee Health And Safety Information Regulatory	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at: <u>https://www.bio-rad-antibodies.com/SDS/MCA1967</u> 10040 For research purposes only				

# Related Products

### **Recommended Secondary Antibodies**

Goat Anti Rat IgG (STAR69)	FITC
Goat Anti Rat IgG (STAR73)	RPE
Rabbit Anti Rat IgG (STAR17)	<u>FITC</u>
Goat Anti Rat IgG (STAR72)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71	DyLight®550, DyLight®650, DyLight®800
Rabbit Anti Rat IgG (STAR21)	HRP
Rabbit Anti Rat IgG (STAR16)	DyLight®800
Goat Anti Rat IgG (STAR131)	Alk. Phos., Biotin
Recommended Useful Reagents	

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & Sout	h Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21	То
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	find a
	Email: antibody_sales_us@	Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-r	ad.com	
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets						
'M415528:230105'						

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint