

## Datasheet: MCA1967

**BATCH NUMBER 172794**

|                      |                       |
|----------------------|-----------------------|
| <b>Description:</b>  | RAT ANTI MOUSE CD44v6 |
| <b>Specificity:</b>  | CD44v6                |
| <b>Format:</b>       | Purified              |
| <b>Product Type:</b> | Monoclonal Antibody   |
| <b>Clone:</b>        | 9A4                   |
| <b>Isotype:</b>      | IgG1                  |
| <b>Quantity:</b>     | 0.1 mg                |

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

|                                | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry                 |     |    | ▪              |                    |
| Immunohistology - Frozen       | ▪   |    |                |                    |
| Immunohistology - Paraffin (1) | ▪   |    |                |                    |
| ELISA                          | ▪   |    |                |                    |
| Immunoprecipitation            |     |    | ▪              |                    |
| Western Blotting               | ▪   |    |                |                    |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.**

|                                 |   |
|---------------------------------|---|
| <b>Target Species</b>           | Mouse   |
| <b>Product Form</b>             | Purified IgG - liquid                             |
| <b>Preparation</b>              | Antibody purified from tissue culture supernatant |
| <b>Buffer Solution</b>          | Phosphate buffered saline                         |
| <b>Preservative Stabilisers</b> | <0.1% sodium azide (NaN <sub>3</sub> )            |

|                                       |  |
|---------------------------------------|--|
| <b>Approx. Protein Concentrations</b> | IgG concentration 0.5mg/ml   |
| <b>Immunogen</b>                      | GST-CD44v6 fusion protein.   |
| <b>External Database Links</b>        | <p><b>UniProt:</b><br/> <a href="#">P15379</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b><br/> <a href="#">12505</a> Cd44    <a href="#">Related reagents</a></p>  |
| <b>Synonyms</b>                       | Ly-24  |
| <b>RRID</b>                           | AB_323213  |
| <b>Fusion Partners</b>                | Spleen cells from immunized female DA rats were fused with cells from the SP/0 myeloma.  |
| <b>Specificity</b>                    | <p><b>Rat anti Mouse CD44v6, clone 9A4</b> recognizes an epitope encoded by exon v6 on the variant portion of murine CD44. CD44v6 has been identified along with ICAM-1 as a co-receptor for Met and in mammary epithelia appears to act at the primary Met co-receptor (<a href="#">Di-Cicco et al. 2015</a>).</p> <p>Rat anti Mouse CD44v6, clone 9A4 has been used successfully for the identification of CD44v6 expressing cells by immunofluorescence (<a href="#">Di-Cicco et al. 2015</a>), western blotting (<a href="#">Lo et al. 2012</a>). Immunohistochemistry (<a href="#">Weilenga et al. 1999</a>) along with ELISA and flow cytometry (<a href="#">Weiss et al. 1997</a>).</p>   |
| <b>References</b>                     | <ol style="list-style-type: none"> <li>Weiss JM <i>et al.</i> (1997) An essential role for CD44 variant isoforms in epidermal Langerhans cell and blood dendritic cell function. <a href="#">J Cell Biol. 137 (5): 1137-47.</a></li> <li>Wielenga VJ <i>et al.</i> (1999) Expression of CD44 in Apc and Tcf mutant mice implies regulation by the WNT pathway. <a href="#">Am J Pathol. 154 (2): 515-23.</a></li> <li>Katagiri, Y.U. <i>et al.</i> (1999) CD44 variants but not CD44s cooperate with beta1-containing integrins to permit cells to bind to osteopontin independently of arginine-glycine-aspartic acid, thereby stimulating cell motility and chemotaxis. <a href="#">Cancer Res. 59: 219-26.</a></li> <li>Hebbard, L. <i>et al.</i> (2000) CD44 expression and regulation during mammary gland development and function. <a href="#">J Cell Sci. 113: 2619-30.</a></li> <li>Rochman, M. <i>et al.</i> (2000) The CD44 receptor of lymphoma cells: structure-function relationships and mechanism of activation. <a href="#">Cell Adhes Commun. 7: 331-47.</a></li> <li>Khaldoyanidi, S. <i>et al.</i> (2002) CD44 variant-specific antibodies trigger hemopoiesis by selective release of cytokines from bone marrow macrophages. <a href="#">Blood. 99: 3955-61.</a></li> <li>Termeer, C. <i>et al.</i> (2003) Targeting dendritic cells with CD44 monoclonal antibodies selectively inhibits the proliferation of naive CD4+ T-helper cells by induction of FAS-independent T-cell apoptosis. <a href="#">Immunology. 109: 32-40.</a></li> <li>Matrosova, V.Y. <i>et al.</i> (2004) Hyaluronic acid facilitates the recovery of hematopoiesis following 5-fluorouracil administration. <a href="#">Stem Cells. 22: 544-55.</a></li> </ol> |

9. Serobyany, N. *et al.* (2005) Nicotinic acetylcholine receptor-mediated stimulation of endothelial cells results in the arrest of haematopoietic progenitor cells on endothelium. [Br J Haematol. 129: 257-65.](#)
10. Boon, E.M. *et al.* (2006) Activation of Wnt signaling in the intestinal mucosa of Apc +/-min mice does not cause overexpression of the receptor tyrosine kinase Met. [Cancer Sci. 97: 710-5.](#)
11. Wallach-Dayyan, S.B. *et al.* (2008) DNA vaccination with CD44 variant isoform reduces mammary tumor local growth and lung metastasis. [Mol Cancer Ther. 7: 1615-23.](#)
12. Lo, B. *et al.* (2012) Lkb1 regulates organogenesis and early oncogenesis along AMPK-dependent and -independent pathways. [J Cell Biol. 199: 1117-30.](#)
13. Nakagawa, H. *et al.* (2014) Loss of liver E-cadherin induces sclerosing cholangitis and promotes carcinogenesis. [Proc Natl Acad Sci U S A. 111\(3\):1090-5.](#)
14. Di-Cicco, A. *et al.* (2015) Paracrine Met signaling triggers epithelial-mesenchymal transition in mammary luminal progenitors, affecting their fate. [Elife. 2015 Jul 13;4.](#)
15. Shibata, W. *et al.* (2017) *Helicobacter*-induced gastric inflammation alters the properties of gastric tissue stem/progenitor cells. [BMC Gastroenterol. 17 \(1\): 145.](#)

---

|                |   |
|----------------|---|
| <b>Storage</b> | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |
|                | Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.   |

---

|                                      |  |
|--------------------------------------|--|
| <b>Guarantee</b>                     | 12 months from date of despatch  |
| <b>Health And Safety Information</b> | Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1967">https://www.bio-rad-antibodies.com/SDS/MCA1967</a> |
| <b>Regulatory</b>                    | For research purposes only   |

---

## Related Products

### Recommended Secondary Antibodies

|  |   |
|--|---|
| Rabbit Anti Rat IgG (STAR16...)                | <a href="#">DyLight@800</a>   |
| Rabbit Anti Rat IgG (STAR17...)                | <a href="#">FITC</a>  |
| Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) | <a href="#">DyLight@550</a> , <a href="#">DyLight@650</a> , <a href="#">DyLight@800</a> |
| Goat Anti Rat IgG (STAR69...)                  | <a href="#">FITC</a>  |
| Goat Anti Rat IgG (STAR73...)                  | <a href="#">RPE</a>   |
| Goat Anti Rat IgG (STAR72...)                  | <a href="#">HRP</a>   |
| Goat Anti Rat IgG (STAR131...)                 | <a href="#">Alk. Phos.</a> , <a href="#">Biotin</a>                                     |
| Rabbit Anti Rat IgG (STAR21...)                | <a href="#">HRP</a>   |

### Recommended Useful Reagents

[ANTIGEN RETRIEVAL BUFFER, pH8.0 \(BUF025A\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M415528:230105'

**Printed on 20 Apr 2026**

---

© 2026 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)