# Datasheet: MCA1969 BATCH NUMBER 161009

Description:	MOUSE ANTI HUMAN CD62E
Specificity:	CD62E
Other names:	E-SELECTIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CL2/6
Isotype:	lgG2a
Quantity:	0.1 mg

## **Product Details**

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				1/20
Immunohistology - Frozen				0.5ug/ml - 2.0ug/ml
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation			•	
Western Blotting			•	
Immunofluorescence	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN <sub>3</sub> )
Approx. Protein Concentrations	IgG concentration 1.0mg/ml

External Database Links	UniProt:			
	P16581 Related reagents			
	Entrez Gene:			
	6401 SELE Related reagents			
Synonyms	ELAM1			
RRID	AB_2186827			
Specificity	<b>Mouse anti Human CD62E antibody, clone CL2/6</b> recognizes human CD62E, a member of the selectin family which is expressed on endothelial cells at sites of inflammation.			
	Mouse anti Human CD62E antibody, clone CL2/6 has been shown to inhibit the binding of human neutrophils to stimulated HUVECS.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.			
References	<ul> <li>Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.</li> <li>1. Abbassi, O. <i>et al.</i> (1993) E-selectin supports neutrophil rolling <i>in vitro</i> under conditions of flow. J Clin Invest. 92 (6): 2719-30.</li> <li>2. Kishimoto, T.K. <i>et al.</i> (1991) Antibodies against human neutrophil LECAM-1 (LAM-1/Leu-8/DREG-56 antigen) and endothelial cell ELAM-1 inhibit a common CD18-independent adhesion pathway <i>in vitro</i>. Blood. 78: 805-11.</li> <li>3. Marin, C. <i>et al.</i> (2011) Mediterranean diet reduces endothelial damage and improves the regenerative capacity of endothelium. Am J Clin Nutr. 93: 267-74.</li> <li>4. Rius, C. <i>et al.</i> (2010) Trans- but not cis-resveratrol impairs angiotensin-II-mediated vascular inflammation through inhibition of NF-kB activation and peroxisome proliferator-activated receptor-gamma upregulation. J Immunol. 185: 3718-27.</li> <li>5. Wassmer, S.C. <i>et al.</i> (2011) Vascular endothelial cells cultured from patients with cerebral or uncomplicated malaria exhibit differential reactivity to TNF. Cell Microbiol. 13: 198-209.</li> <li>6. Nieuwland, R. <i>et al.</i> (2000) Cellular origin and procoagulant properties of microparticles in meningococcal sepsis. Blood. 95: 930-5.</li> <li>7. Lo. C. Y. <i>et al.</i> (2013) The use of surface immobilization of P-selectin glycoprotein ligand-1 on mesenchymal stem cells to facilitate selectin mediated cell tethering and rolling <u>Biomaterials. 34 (33): 8213-22.</u></li> </ul>			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.			
Guarantee	12 months from date of despatch			
Health And Safety	Material Safety Datasheet documentation #10040 available at:			

Information	https://www.bio-rad-antibodies.com/SDS/MCA1969 10040			
Regulatory	For research purposes only			

### **Related Products**

#### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>			
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos., DyLight®488, DyLight®550,</u>			
	<u>DyLight®650, DyLight®680, DyLight®800,</u>			
	FITC, HRP			
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>			
Goat Anti Mouse IgG (STAR77)	HRP			
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP			
<b>Recommended Negative Controls</b>				

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389355:210806'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint