

Datasheet: MCA1969F BATCH NUMBER 166696

Description:MOUSE ANTI HUMAN CD62E:FITSpecificity:CD62EOther names:E-SELECTINFormat:FITCProduct Type:Monoclonal AntibodyClone:CL2/6Isotype:IgG2aQuantity:100 TESTS		
Other names:E-SELECTINFormat:FITCProduct Type:Monoclonal AntibodyClone:CL2/6Isotype:IgG2a	Description:	MOUSE ANTI HUMAN CD62E:FITC
Format:FITCProduct Type:Monoclonal AntibodyClone:CL2/6Isotype:IgG2a	Specificity:	CD62E
Product Type: Monoclonal Antibody Clone: CL2/6 Isotype: IgG2a	Other names:	E-SELECTIN
Clone: CL2/6 Isotype: IgG2a	Format:	FITC
Isotype: IgG2a	Product Type:	Monoclonal Antibody
	Clone:	CL2/6
Quantity: 100 TESTS	lsotype:	lgG2a
-	Quantity:	100 TESTS

Product Details

Applications	derived from testing w communications from	ithin our laboratories the originators. Pleas al protocol recomme	he following application , peer-reviewed publica se refer to references in ndations, please visit <u>w</u>	tions or personal dicated for further
		Yes No	Not Determined	Suggested Dilution
	Flow Cytometry	•		
	necessarily exclude its	s use in such proced mmended that the us	er titrates the product f	g dilutions are given as
Target Species	Human			
Product Form	Purified IgG conjugate	d to Fluorescein Isot	hiocyanate Isomer 1 (F	ITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	
Preparation	Antibody purified from	tissue culture super	natant	
Buffer Solution	TRIS buffered saline			
Preservative Stabilisers	0.02% sodium azide (l 1% bovine serum albu	0,		
External Database Links	UniProt:			

	P16581 Related reagents
	Entrez Gene:
	6401 SELE Related reagents
Synonyms	ELAM1
RRID	AB_323219
Specificity	Mouse anti Human CD62E antibody, clone CL2/6 recognizes human CD62E, a member of the selectin family which is expressed on endothelial cells at sites of inflammation.
	Mouse anti Human CD62E antibody, clone CL2/6 has been shown to inhibit the binding of human neutrophils to stimulated HUVECS.
Flow Cytometry	Use 10μ I of the suggested working dilution to label 10^6 cells in 100μ I
References	 Kishimoto, T.K. <i>et al.</i> (1991) Antibodies against human neutrophil LECAM-1 (LAM-1/Leu-8/DREG-56 antigen) and endothelial cell ELAM-1 inhibit a common CD18-independent adhesion pathway <i>in vitro</i>. <u>Blood. 78: 805-11.</u> Abbassi, O. <i>et al.</i> (1993) E-selectin supports neutrophil rolling <i>in vitro</i> under conditions of flow. <u>J Clin Invest. 92 (6): 2719-30.</u> Nieuwland, R. <i>et al.</i> (2000) Cellular origin and procoagulant properties of microparticles in meningococcal sepsis. <u>Blood. 95: 930-5.</u> Rius, C. <i>et al.</i> (2010) Trans- but not cis-resveratrol impairs angiotensin-II-mediated vascular inflammation through inhibition of NF-kB activation and peroxisome proliferator- activated receptor-gamma upregulation. <u>J Immunol. 185: 3718-27.</u> Marin, C. <i>et al.</i> (2011) Mediterranean diet reduces endothelial damage and improves the regenerative capacity of endothelium. <u>Am J Clin Nutr. 93: 267-74.</u> Wassmer, S.C. <i>et al.</i> (2011) Vascular endothelial cells cultured from patients with cerebral or uncomplicated malaria exhibit differential reactivity to TNF. <u>Cell Microbiol. 13: 198-209.</u> Lo. C. Y. <i>et al.</i> (2013) The use of surface immobilization of P-selectin glycoprotein ligand-1 on mesenchymal stem cells to facilitate selectin mediated cell tethering and rolling <u>Biomaterials. 34 (33): 8213-22.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10058 available at: https://www.bio-rad-antibodies.com/SDS/MCA1969F

	10058				
Regulato	ry For res	earch purpose	s only		
Relate	d Products				
Recomn	nended Negative C	Controls			
MOUSE lo	G2a NEGATIVE CONT	ROL:FITC (MCA	<u> 4929F)</u>		
Recomn	nended Useful Rea	agents			
	EROBLOCK (BUF070A				
<u>HUMAN S</u>	EROBLOCK (BUF070E	<u>3)</u>			
rth & South herica	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
leneu	Email: antibody sales us@b	io-rad.com	Email: antibody_sales_uk@bic	-rad.com	Email: antibody_sales_de@bio-rad.co

'M423280:231011'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint