

Datasheet: MCA1971GA

BATCH NUMBER 162371

Description:	MOUSE ANTI PIG CD16
Specificity:	CD16
Other names:	FcRIII
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	G7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Porcine peripheral blood leucocytes
External Database Links	<p>UniProt: Q28942 Related reagents</p> <p>Entrez Gene: 397684 FCGR3B Related reagents</p>
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the mouse P3-X63-Ag8.653 myeloma cell line
Specificity	<p>Mouse anti Pig CD16, clone G7 recognizes porcine CD16 also known as Fc-gamma RIII or the low affinity IgG (Fc) receptor III. Clone G7 was clustered as CD16 at the Second International Workshop to Define Swine Cluster of Differentiation (CD) Antigens (Saalmuller et al. 1998).</p> <p>Mouse anti pig CD16 immunoprecipitates a protein of ~40 kDa from porcine neutrophils and NK cells (Wierda et al. 1993). Subsequent cloning and characterization of the G7 molecule indicated that G7 was the porcine homologue of Human CD16 (Halloran et al. 1994).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Dato, M.E. <i>et al.</i> (1992) A triggering structure recognized by G7 monoclonal antibody on porcine lymphocytes and granulocytes. Cell Immunol. 140 (2): 468-77. Wierda, W.G. <i>et al.</i> (1993) Two distinct porcine natural killer lytic trigger molecules as PNK-E/G7 molecular complex. Cell Immunol. 146 (2): 270-83. Halloran, P.J. <i>et al.</i> (1994) Biochemical characterization of the porcine Fc gamma RIII alpha homologue G7. Cell Immunol. 158 (2): 400-13. Devriendt, B. <i>et al.</i> (2010) Targeting of <i>Escherichia coli</i> F4 fimbriae to Fcgamma receptors enhances the maturation of porcine dendritic cells. Vet Immunol Immunopathol. 135 (3-4): 188-98. Inman, C.F. <i>et al.</i> (2010) Dendritic cells interact with CD4 T cells in intestinal mucosa. J Leukoc Biol. 88 (3): 571-8. Terzic, S. <i>et al.</i> (2002) Immunophenotyping of leukocyte subsets in peripheral blood and palatine tonsils of prefattening pigs. Vet Res Commun. 26: 273 - 83. Masure, D. <i>et al.</i> (2013) A Role for Eosinophils in the Intestinal Immunity against Infective <i>Ascaris suum</i> Larvae. PLoS Negl Trop Dis. 7: e2138. Hester, S.N. <i>et al.</i> (2012) Intestinal and systemic immune development and response to vaccination are unaffected by dietary (1,3/1,6)-β-D-glucan supplementation in neonatal piglets. Clin Vaccine Immunol. 19 (9): 1499-508. Kapetanovic, R. <i>et al.</i> (2012) Pig bone marrow-derived macrophages resemble human

- macrophages in their response to bacterial lipopolysaccharide. [J Immunol. 188: 3382 - 94.](#)
10. Gimeno, M. *et al.* (2011) Cytokine profiles and phenotype regulation of antigen presenting cells by genotype-I porcine reproductive and respiratory syndrome virus isolates. [Vet Res. 42: 9.](#)
11. Mussá, T. *et al.* (2011) Interaction of porcine conventional dendritic cells with swine influenza virus. [Virology 420: 125-34.](#)
12. Vincent, I.E. *et al.* (2003) Dendritic cells harbor infectious porcine circovirus type 2 in the absence of apparent cell modulation or replication of the virus. [J Virol. 77: 13288 - 300.](#)
13. Inman, C.F. *et al.* (2012) Neonatal colonisation expands a specific intestinal antigen-presenting cell subset prior to CD4 T-cell expansion, without altering T-cell repertoire. [PLoS One 7: e33707.](#)
14. Sánchez, C. *et al.* (1999) The porcine 2A10 antigen is homologous to human CD163 and related to macrophage differentiation. [J Immunol. 162 \(9\): 5230-7.](#)
15. Lecours, M.P. *et al.* (2011) Characterization of porcine dendritic cell response to *Streptococcus suis*. [Vet Res. 42: 72.](#)
16. Inman, C.F. *et al.* (2010) Rearing environment affects development of the immune system in neonates. [Clin Exp Immunol. 160 \(3\): 431-9.](#)
17. Summerfield, A. *et al.* (2003) Porcine peripheral blood dendritic cells and natural interferon-producing cells. [Immunology 110: 440-9.](#)
18. Mair, K.H. *et al.* (2012) NKp46 expression discriminates porcine NK cells with different functional properties. [Eur J Immunol. 42: 1261-71.](#)
19. Mair, K.H. *et al.* (2013) Porcine CD8 α dim⁻/NKp46^{high} NK cells are in a highly activated state. [Vet Res. 44: 13.](#)
20. Auray, G. *et al.* (2016) Characterization and Transcriptomic Analysis of Porcine Blood Conventional and Plasmacytoid Dendritic Cells Reveals Striking Species-Specific Differences. [J Immunol. Nov 11. pii: 1600672. \[Epub ahead of print\]](#)
21. Kyrova, K. *et al.* (2014) The response of porcine monocyte derived macrophages and dendritic cells to *Salmonella typhimurium* and lipopolysaccharide. [BMC Vet Res. 10: 244.](#)
22. Suzuki, S. *et al.* (2016) Generation and characterization of RAG2 knockout pigs as animal model for severe combined immunodeficiency. [Vet Immunol Immunopathol. 178: 37-49.](#)
23. Waide, E.H. *et al.* (2015) Not All SCID Pigs Are Created Equally: Two Independent Mutations in the Artemis Gene Cause SCID in Pigs. [J Immunol. 195 \(7\): 3171-9.](#)
24. Loss, H. *et al.* (2018) Effects of a pathogenic ETEC strain and a probiotic Enterococcus faecium strain on the inflammasome response in porcine dendritic cells. [Vet Immunol Immunopathol. 203: 78-87.](#)
25. LeLuduec, J.B. *et al.* (2016) Intradermal vaccination with un-adjuvanted sub-unit vaccines triggers skin innate immunity and confers protective respiratory immunity in domestic swine. [Vaccine. 34 \(7\): 914-22.](#)
26. Ferret-Bernard, S. *et al.* (2020) Maternal Supplementation of Food Ingredient (Prebiotic) or Food Contaminant (Mycotoxin) Influences Mucosal Immune System in Piglets. [Nutrients. 12 \(7\): 2115.](#)
27. Skovdal, S.M. *et al.* (2019) Inhaled nebulized glatiramer acetate against Gram-negative bacteria is not associated with adverse pulmonary reactions in healthy, young adult female pigs. [PLoS One. 14 \(10\): e0223647.](#)
28. Fernández-Caballero, T. *et al.* (2018) Phenotypic and functional characterization of

- porcine bone marrow monocyte subsets. [Dev Comp Immunol. 81: 95-104.](#)
29. Teuben, M.P.J. *et al.* (2021) Standardized porcine unilateral femoral nailing is associated with changes in PMN activation status, rather than aberrant systemic PMN prevalence. [Eur J Trauma Emerg Surg. Jun 10 \[Epub ahead of print\].](#)
30. Teuben, M. *et al.* (2021) Instant intra-operative neutropenia despite the emergence of banded (CD16^{dim}/CD62L^{bright}) neutrophils in peripheral blood - An observational study during extensive trauma-surgery in pigs. [Injury. 52 \(3\): 426-33.](#)
31. Van der Weken, H. *et al.* (2021) Antibody-Mediated Targeting of Antigens to Intestinal Aminopeptidase N Elicits Gut IgA Responses in Pigs. [Front Immunol. 12: 753371.](#)
32. Boettcher, A.N. *et al.* (2020) CD3ε⁺ Cells in Pigs With Severe Combined Immunodeficiency Due to Defects in ARTEMIS [Frontiers in Immunology. 11 \[Epub ahead of print\].](#)
33. Zhao, H. *et al.* (2022) Development of *RAG2*^{-/-} *IL2Rγ*^{-/-} immune deficient FAH-knockout miniature pig. [Front Immunol. 13: 950194.](#)

Further Reading	<p>1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.</p> <p>2. Gerner W <i>et al.</i> (2015) Phenotypic and functional differentiation of porcine αβ T cells: current knowledge and available tools. Mol Immunol. 66 (1): 3-13.</p>
------------------------	--

Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p>
----------------	--

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	<p>Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1971GA 10040</p>
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)

Rabbit Anti Mouse IgG (STAR9...) [FITC](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M383311:210513'

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)